arXiv:1108.4203v3 [astro-ph.CO] 1 Feb 2012

Signatures of very high energy physics in the squeezed limit of the bispectrum from the field theoretical approach

Diego Chialva
Université de Mons, Service de Mécanique et gravitation, Place du Parc 20, 7000 Mons, Belgium
`diego.chialva@umons.ac.be`

Abstract

We investigate the signatures, in the squeezing limit of the primordial scalar bispectrum, of modifications of the standard theory at high energy, such as a modified quantum vacuum or modified dispersion relations, reflecting the effects of different physics at higher energies/earlier times. Our results are obtained through the analysis in field theory and not using the (approximated) templates for the bispectrum. In this way we can obtain the precise dependence on the probed scale and quantify the presence of enhancements and/or non-local shape of the non-Gaussianities, which are relevant, for example, for measurements of the halo bias. It is shown that the signatures are distinctive and can be determined in general, and the results are different from those obtained using the templates proposed in the literature, where existing, in particular because of the interplay between the scale of the observations and the scale of the high-energy physics when taking the large-scale (squeezed) limit. Several pieces of information regarding high energy physics could be obtained in case of detection of these signals, especially bounds on the scales of new physics.

Contents

1 Introduction **1**

2 Formalism and notation. **3**

3 Brief review of the scenarios under consideration. **4**

4 Bispectrum in the squeezing limit **8**

5 Signatures of very high energy physics in the squeezed limit **21**

6 Comments on the halo bias **26**

7 Final discussion and conclusion **27**

A Appendices **28**

1 Introduction

The non-Gaussian features of the primordial cosmological perturbations are going to become one of the most important experimental and theoretical tools for investigating the early history of the Universe [1]. The scalar bispectrum, coming from the three-point correlator of the comoving curvature perturbation ζ, is the leading contribution to the scalar non-Gaussianities in the perturbative theory. In recent times, several reasons of interests have emerged for the investigation of its so-called *squeezed limit*, when one of the three momenta it depends on in Fourier space is much smaller than the other two: $k_1 \ll k_S, k_{2,3} \simeq k_S$.

In fact, large-scale probes of the primordial non-Gaussianities like the halo bias are expected to be soon competing with or even surpass in accuracy those of the CMBR [2, 3], and this prompts for a better understanding of their sensitivity on the details of the bispectrum and the theoretical models.

Furthermore, studying the squeezed limit provides powerful results capable of distinguishing (hopefully falsify) classes of models. For example, all single-field models (adiabatic perturbations) with Bunch-Davies vacuum and standard kinetic terms (we call this the standard scenario from now on) predict the so-called *local* form of the bispectrum and a very low level of non-Gaussianities in the limit [4–6]. Remarkably, measurements of the halo bias could discriminate cases with different underlying assumptions [7, 8]: for example models with modified kinetic terms whose bispectrum template is maximized on *equilateral* configurations, or with initial state different than the Bunch-Davies, in which case the proposed template is maximized on *folded* configurations [9].

In all the analysis in the literature, however, the halo bias and the squeezing limit have been investigated using the *templates* and not the actual field theoretical results for the bispectrum [1]. Sometimes the discrepancies between templates and field theoretical results disappear in the squeezing limit: for example, Maldacena's result [4] becomes factorizable in that limit, and assumes indeed a local shape. However, we will show that in other cases the use of templates in place of the actual theoretical predictions appreciably changes the outcomes.

---

[1] While completing this paper, we have found that in [10] the analysis and the simulations regarding the halo bias are still done using the templates, but the authors warn that they could lead to the wrong predictions for the observables. However, the detailed study of the field theoretical results was not done, in particular not for the folded template. After the appearance of this article, we were also signalled [11], which also makes some remarks, focusing on the presence of enhancements, about the squeezing limit in a particular formalism (density matrix approach) with a specific initial condition for the field mode functions. However, also in this case the detailed study of the general field theoretical results was not undertaken (the conclusions in that paper are in agreement with those in one of the classes of scenarios we deal with).

In this paper, we want to look at the field theory cosmological predictions for the squeezed limit of the bispectrum for single-field slow-roll inflationary models, investigating the effects of two very simple modifications of the standard scenario: 1) the presence of higher derivative corrections to the action that are unsuppressed at a certain physical scale Λ and modify the dispersion relations of the perturbation fields [12], and 2) the choice of an initial state other than the Bunch-Davies vacuum, reflecting the effects of different physics at higher energies/earlier times as proposed in [13, 14].

The description of the physics at high energies is largely unknown, and, at the same time, there are various different models of single-field inflation. It is therefore important to derive general results, as indeed was in the spirit of [5, 6]. We will therefore deal with this topic adopting the same phenomenological approach of [9, 15, 16], not assuming specific detailed high-energy models, but still obtaining quantitative as well as qualitative results. We will present in more details the approach and the scenarios in section 3. It is also important to stress that we consider the realistic squeezing limit, where k_1 is small but non-zero, as defined in [6–8], in view of possible future observations.

A part from [9,15], examples of the modification to the vacuum that we describe have been proposed in [13, 14] and many successive papers. On the other hand, modified dispersion relations occur in a variety of models, such as certain realization of quantum gravity, as the Hořava-Lifshitz model [17], or also in braneworld models [18], and are of interest for the studies of Lorentz violation [19]. They occur as well in the effective theory when the fields propagate with small speed of sound [20].

Before presenting the detailed analysis, it is useful to outline the points that could make the difference with the standard scenario in the cases we consider. In fact, the standard result follows from a simple argument [4–6]: in the standard scenario, the leading contribution to the non-Gaussianities comes from late times, and furthermore, in the squeezing limit, we consider one of the three momenta (k_1) that the bispectrum depends upon to be very small. Therefore the leading contribution to non-Gaussianities occurs when $k_1\eta \ll 1$ (η is the conformal time), and thus the perturbation depending on k_1 is at superhorizon scales, is frozen and acts as a background for the other perturbations in the bispectrum. It has the effect of shifting the horizon-crossing time for the other perturbations, and the result is thus determined by the (slow-roll suppressed) scalar tilt of the spectrum, see [4–6]. It has been argued that, because of the generality of this argument, this result for the squeezing limit of the bispectrum is valid for any single-field slow-roll model –with Bunch-Davies vacuum and standard dispersion relations–, with all possible couplings that allow a slow-roll inflation.

However, this argument would fail for models with an additional important contribution to the non-Gaussianities at much earlier times when it still is $k_1\eta > 1$ and the perturbation is not frozen at superhorizon scales. The scenarios of modified initial vacuum and modified dispersion relations have in fact such additional early time contribution to the non-Gaussianities: it is due to particle creation, respectively at the time when the condition picking up the non-Bunch-Davies vacuum is imposed, and at the time/scale of appearance of the new physics that modifies the dispersion relations. One could think that the particle creation would represent a tiny effect, being severely constrained by backreaction (as we will review), but the bispectrum can be quite sensitive to it, because particle creation will lead also to interference effect and phase cancellations in the correlators that could lead to enhancements.

We want to investigate the effects of these contributions on the squeezing limit of the bispectrum by performing the analysis at the rigorous level of the field theory description, investigating whether general features appear and if violations of the standard result occur. It is important to make two comments regarding our work. First, the analysis we make is general, within the scenarios of section 3, and ranges over all possible single-field slow roll models, allowing all those operators that permit a slow-roll inflationary evolution. Although many operators would be quite constrained or suppressed because of backreaction/preservation of slow-roll, as it was discussed in [6,21], the allowed ones are still numerous, and one could think that looking for the emergence of general features would be hopeless.

However, the squeezing limit allows powerful results that rely on very general features of the scenarios, common to all the specific models, permitting, as we said before, to constrain entire classes of models. In the case of the standard scenario, the constraints following from the freeze-out argument we reviewed above are so strong that the full leading form of the squeezing limit of the bispectrum is obtained, valid for any model of that scenario, while the differences among the specific models appear only in the subleading (by powers of $\frac{k_1}{k_S}$) corrections. Also in the modified scenarios we are considering, there are general constraining arguments (particle creation, interference phenomena), and we will show that the squeezing limit is in fact dominated by these general aspects of the scenarios, at leading order, and thus there appear general features in the squeezed bispectrum.

Second, let us recall that in the case of modified initial vacuum, one can divide the contributions to non-Gaussianities in two parts [9, 15]: one deriving from intrinsic non-Gaussianities at the boundary where the boundary condition defining the initial state is imposed, the other from the particle production due to the modified vacuum. The first kind of contribution is absent for Gaussian initial state modifications. As in [9, 13, 15], we focus on this class of modifications, which represents the minimal deviation from the Bunch-Davies vacuum. We will show that already this minimal change of initial conditions has quite an effect on the squeezing limit. Moreover, at least in the boundary action formalism, the leading intrinsic non-Gaussian modifications have been shown to be of the local type –see [22] and the comment in [9]– and, being also quite constrained by backreaction, they would thus lead to a squeezing limit similar to that in the standard scenario.

In the case of modifications of the initial quantum state, we will also show that the field theoretic result in the squeezing limit is very different from those obtained in previous studies, see [7, 8], using the folded template proposed for CMBR analysis [9]. This disagreement could have been already anticipated, since the standard evaluators (cosine and fudge factor) for the matching between the proposed template [9] and the theoretical prediction [15] generically indicate that the two depart more and more for large $k_L\eta_c$, where k_L is the largest momentum and η_c is the time when the boundary condition picking up the non-Bunch-Davies vacuum is imposed, see [9]. Even more importantly, the template does not depend on the parameter η_c, and thus taking the squeezing limit in the full result might (and will) be different that doing it in the template, because of the presence of distinct scales.

The outline of the paper is as follows: we start in section 2 with a very rapid presentation of the necessary formalism and notation, followed by a brief review in section 3 of the scenarios that we deal with. The analysis of the realistic squeezed limit of the bispectrum is reported in sections 4 and 5. First, we present in sections 4.1 and 4.2 the detailed computations for two examples of interactions (which will be also proven to yield among the largest contributions to the bispectrum), to illustrate step-by-step how their contributions are affected by the features of the modified scenarios. Then, we generalize the study to the other various possible couplings in section 4.3. The analysis and discussion on the results are contained in section 5. Finally, we briefly comment on the consequences of our results on the halo bias in section 6, and conclude in 7. Some useful equations and material, which would have made heavy the main text, have been moved to the appendices.

2 Formalism and notation.

We begin by introducing the basic notation and elements of the formalism of cosmological perturbation theory, for more details see [4]. Scalar perturbations in single-field slow-roll inflationary models are efficiently parametrized by the comoving curvature perturbation[2]

$$\zeta(\eta, \vec{x}) = \int \frac{d^3k}{(2\pi)^{\frac{3}{2}}} \zeta_{\vec{k}}(\eta) e^{i\vec{k}\cdot\vec{x}}. \tag{1}$$

---

[2] We follow the conventions in [4, 15, 16]; other authors call this perturbation $\mathcal{R}$ to distinguish it from the uniform energy-density curvature perturbation.

The slow-roll parameters are defined as $\epsilon = \frac{\dot\phi^2}{2H^2M_{\text{Planck}}^2}$, $\eta_{\text{sl}} = -\frac{\ddot\phi}{\dot\phi H} + \frac{\dot\phi^2}{2H^2M_{\text{Planck}}^2}$, H is the Hubble rate. ϕ the inflaton (background), and $a(\eta) \sim -\frac{1}{H\eta}$ is the scale factor at zeroth order in slow-roll. Throughout the paper, dots (primes) indicate derivatives with respect to cosmic time t (conformal time η).

The two-point function is defined as

$$\langle \zeta(\vec{k}_1)\zeta(\vec{k}_2)\rangle = (2\pi)^3\delta^{(3)}(\vec{k}_1+\vec{k}_2)\,P(k), \qquad k = |\vec{k}_1|\,. \tag{2}$$

and, following [23], we define the bispectrum B as

$$\langle \zeta_{\vec{k}_1}(\eta)\zeta_{\vec{k}_2}(\eta)\zeta_{\vec{k}_3}(\eta)\rangle = (2\pi)^3\delta(\sum_i \vec{k}_i)\hat{B}(\vec{k}_1,\vec{k}_2,\vec{k}_3,\eta)\,. \tag{3}$$

A key ingredient in the computation of correlators is the Whightman function

$$(2\pi)^3\delta^{(3)}(\vec{k}_1+\vec{k}_2)\,G_{\vec{k}}(\eta,\eta') \equiv \langle \zeta_{\vec{k}_1}(\eta)\zeta_{\vec{k}_2}(\eta')\rangle = (2\pi)^3\delta^{(3)}(\vec{k}_1+\vec{k}_2)\,\frac{H^2}{\dot\phi^2}\frac{f_k(\eta)}{a(\eta)}\frac{f_k^*(\eta')}{a(\eta')} \tag{4}$$

where $f_{\vec{k}}(\eta), f^*_{\vec{k}}(\eta)$ are two linearly independent solutions of the field equation

$$f''_{\vec{k}} + \left(\omega(\eta,\vec{k})^2 - \frac{z''}{z}\right) f_{\vec{k}} = 0 \qquad z = \frac{a\dot\phi}{H}\,, \tag{5}$$

such that, once quantized,

$$\zeta_{\vec{k}}(\eta) = \frac{f_{\vec{k}}(\eta)}{z}\hat{a}^\dagger_{\vec{k}} + \frac{f^*_{\vec{k}}(\eta)}{z}\hat{a}_{\vec{k}}\,. \tag{6}$$

Imposing the standard commutation relation on the operators $\hat{a}_{\vec{k}}{}^\dagger, \hat{a}_{\vec{k}}$, entails a certain normalization for the Wronskian of $f^*_{\vec{k}}(\eta), f_{\vec{k}}(\eta)$. Different choices for $f^*_{\vec{k}}(\eta), f_{\vec{k}}(\eta)$ are different choices of vacua [4]. In equation (5), ω is the (comoving) frequency as read from the effective action: we will limit ourselves to the isotropic case, where ω depends only on k, dropping the arrow symbol.

3 Brief review of the scenarios under consideration.

To ease the discussion of the results for the three-point function and make the paper self-contained, let us recap here the main features of the scenarios we will deal with, for more details see [9, 12–16].

3.1 The standard scenario

In the standard scenario the dispersion relation is the standard Lorentzian one and the mode functions $f_{\vec{k}}(\eta)$ are determined by choosing the Bunch-Davies vacuum, so that

$$\omega(\eta,k)^2 = k^2\,, \qquad f_k(\eta) = \frac{\sqrt{\pi}}{2}e^{i\frac{\pi}{2}\nu + i\frac{\pi}{4}}\,\sqrt{-\eta}H_\nu^{(1)}(-k\eta)\,, \tag{7}$$

where $H^{(1)}$ is the first Hankel's functions with $\nu = \frac{3}{2} + \frac{1-n_s}{2}$, and the equations and solutions are extrapolated up to $\eta_{\text{in}} = -\infty$ where the initial state has been chosen as the empty vacuum (Bunch-Davies). $n_s = 1 - 6\epsilon + 2\eta_{\text{sl}}$ is the spectral index. For small momenta, at leading order in ϵ,

$$P_{\text{st}}(k) \underset{k\to 0}{\sim} \frac{H^2}{4M_{\text{Planck}}^2\,\epsilon k^3} \tag{8}$$

where the suffix $_{\text{st}}$ indicates that we are considering the standard scenario.



Figure 1: Generic example of dispersion relation with violation of WKB at early times.

3.2 Modified dispersion relations

Higher-derivative corrections to the action are deeply rooted in the well-established effective field theory framework. In fact, one expects that at certain times at least some of the higher derivative corrections to the action cannot be treated perturbatively, because, since the metric scale factor $a(\eta)$ is rapidly decreasing back in time, the suppressing factors, given by powers of ratios such as $\frac{p}{\Lambda} = \frac{k}{a(\eta)\Lambda}$, where Λ is the relevant high-energy scale, are not small any more. These no longer small corrections can lead to modified dispersion relations. As recalled in the introduction, such possibility appears in a variety of scenarios, see for example those in [17–20].

As we said, we adopt a phenomenological approach and consider *generic* modifications to the dispersion relation due to higher-derivative corrections to the effective action, encoding them in a function $F(-\frac{p}{\Lambda})$ by writing the comoving frequency as

$$\omega(\eta, k) = a(\eta)\omega_{\text{phys}}(p) = a(\eta)p\, F\left(-\frac{p}{\Lambda}\right) = k\, F\left(\frac{H}{\Lambda}k\eta\right), \qquad F(x \to 0) \to 1\,, \qquad \frac{H}{\Lambda} \gg 1\,. \tag{9}$$

In [16, 24], it has been shown that if the modified dispersion relations always satisfy the WKB conditions (adiabaticity) at early times (that is, WKB is violated only when the dispersion is in the standard linear regime $\omega \simeq k$ as usual) the bispectrum is similar to that of the standard scenario, and so it will be its squeezing limit. This is easily understood, since the particle production is very limited in those cases.

Instead, [16] has shown that the bispectrum is particularly sensitive to modifications that lead to a violation of the WKB conditions (adiabaticity) at early times, where particle production is more substantial. We will therefore focus on those scenarios.

The generic shape of a dispersion relation with WKB violation once at early times is depicted in figure 1. We stress that we are not proposing any specific model, which would be a strong assumption on the high energy physics: we consider the most generic function $F\left(-\frac{p}{\Lambda}\right)$ in equation (9), with the only constraint that adiabaticity is once violated at early times (it is easy to extend our analysis to multiple violations). We are in fact interested in the general consequences on the bispectrum from this scenario. Similar phenomenological approach and generic dispersion relations have been considered, studying the spectrum, for example in the references [12], see also [19].

A general treatment of the field equation (5) in these cases yields the following solution [16]:

$$f_k(\eta) = \begin{cases} \varsigma_k\, u_1(\eta, k) & \text{I}:\ \eta < \eta_{\text{I}}^{(k)} \\ B_1\, \mathcal{U}_1(\eta, k) + B_2\, \mathcal{U}_2(\eta, k) & \text{II}:\ \eta_{\text{I}}^{(k)} < \eta < \eta_{\text{II}}^{(k)} \\ \alpha_k^{\text{mdr}}\, u_1(\eta, k) + \beta_k^{\text{mdr}}\, u_2(\eta, k) & \text{III}:\ \eta_{\text{II}}^{(k)} < \eta < \eta_{\text{III}}^{(k)} \\ D_1\, \mathcal{V}_1(\eta, k) + D_2\, \mathcal{V}_2(\eta, k) & \text{IV}:\ \eta_{\text{III}}^{(k)} < \eta \end{cases} \tag{10}$$

where $u_{1,2}(\eta, k)$ can be very well approximated using the WKB method, while $\mathcal{V}_{1,2}(\eta, k)$ and $\mathcal{U}_{1,2}(\eta, k)$ need to be found by other means, as the WKB conditions are not satisfied [3]. As known, given equation (5), the times $\eta^{(k)}_{\text{I},\text{II},\text{III}}$ where the WKB approximation first fails are in proximity of the turning points of $V(\eta, k)^2 \equiv \omega(\eta, k)^2 - \frac{z''}{z}$, see [25, 26]. They depend on k; however, when clear from the context, we will omit the label $^{(k)}$. The coefficients $D_{1,2}, B_{1,2}, \alpha_k^{\text{mdr}}, \beta_k^{\text{mdr}}$ are obtained by asking for the continuity of the function and its first derivative, and by imposing the Wronskian condition $\mathcal{W}\{f, f^*\} = -i$ in order to have the standard commutation relations in the quantum theory.

The details of the solution can be found in [16], and are briefly reviewed in the appendix A.1. Some of the salient features needed here are that:

- backreaction constraints the interval $[\eta_{\text{I}}, \eta_{\text{II}}]$ of WKB violation to be very small, that is
$$\Delta = \frac{\eta_{\text{I}} - \eta_{\text{II}}}{\eta_{\text{I}}} \ll 1\,, \tag{11}$$
- expanding for small Δ, we obtain generically (i.e., independently of the detailed form of $\omega(\eta, k)$)
$$\alpha_k^{\text{mdr}} = \left(1 - i\frac{V(\eta_{\text{I}}, k)\,\eta_{\text{I}}}{2}\frac{\mathcal{Q}}{V^2}\bigg|_{\eta_{\text{I}}}\Delta + \mathcal{O}(\Delta^2)\right)\varsigma_k\,, \qquad \beta_k^{\text{mdr}} = \left(i\frac{V(\eta_{\text{I}}, k)\,\eta_{\text{I}}}{2}\frac{\mathcal{Q}}{V^2}\bigg|_{\eta_{\text{I}}}\Delta + \mathcal{O}(\Delta^2)\right)e^{-\frac{2i}{\epsilon}\Omega|_{\eta_{\text{I}}}}\varsigma_k\,, \tag{12}$$
where $V(\eta, k)^2 \equiv \omega(\eta, k)^2 - \frac{z''}{z}$. Here, $\frac{\mathcal{Q}}{V^2}\big|_{\eta_{\text{I}}}$ is the order 1 factor signalling the violation of the WKB conditions at η_{I}, and $\mathcal{Q}$ is reported in equation (100) in appendix A.1. The magnitude of $|\beta_k^{\text{mdr}}|$ is constrained by backreaction. We will discuss more of the k-dependence and the magnitude of these coefficients in section 3.4,
- we choose $\varsigma_k = 1$ picking up the usual adiabatic vacuum.

The two-point function in the small momentum limit, at leading order in ϵ, $\beta_k^{\text{mdr}}, \Delta$, is
$$P_{\text{mdr}}(k) \underset{k\to 0}{\sim} \frac{H^2}{4M^2_{\text{Planck}}\,\epsilon k^3}\left(1 + 2\,\text{Re}(\beta_k^{\text{mdr}})\right). \tag{13}$$

3.3 Modified initial state

Let us review the two implementations of the modified vacuum (initial state) approach that have been first proposed in [13, 14], reflecting the effects of different physics at higher energies/earlier times than inflation in setting the initial/boundary condition for the perturbation fields:

- EBA approach: fix the boundary conditions for the fields at the same specific finite time η_c (beginning of inflation) independently of the modes k, through an effective boundary action (EBA) which accounts for the high energy physics via renormalization [14]. This kind of boundary condition strongly breaks scale invariance, being imposed for all modes k at the same boundary time. The (cutoff) scale Λ, proper of the effective action formulation, is not related to the boundary time η_c.
- NPHS approach: consider the effective theory valid up to a certain energy scale Λ, and choose the boundary conditions for the solution to the field equation at the new physics hyper-surface (NPHS) corresponding to when the physical momentum reaches that cutoff scale [13]. In this case, the cutoff is imposed in a scale invariant way via the condition $\frac{k}{a(\eta_c)} = \Lambda$, so that the time η_c when the initial state is picked is k- and Λ- dependent. At zero slow-roll order, $\eta_c = -\frac{\Lambda}{kH}$.

---

[3] Some different general approximations are possible, see [16] and the appendix A.1. With particular choices of time coordinate, also the WKB approximation can be applied sometimes, see [25].

To spare notation, we use the symbol η_c for both the NPHS and EBA cases and make clear in the context which approach we are following.

As in [9, 15], when studying these scenarios we further adopt a phenomenological method and consider generic Gaussian boundary conditions. The latter is a minimal change from the Bunch-Davies vacuum. In this case the three- and higher- point functions are determined by the two-point function. However, recall that in principle the boundary condition defining the modified vacuum can be chosen to be intrinsically non-Gaussian, leading to further contributions to the three-point function not determined by the two-point one. Let us also point out that, in fact, in the EBA language, it was shown in [22] –see also the comment in [9]–, that the leading corrections due to intrinsic non-Gaussianities in the boundary condition would be of the local form, which would thus yield the same form of squeezing limit as in the standard scenario (with a suppressed amplitude, because of backreaction and lack of cumulation with time, see later sections). We will however focus on the minimal Gaussian vacuum modification, as done in [9, 15] (see also the study of the related observational prediction's in [7–9])

Then, both in the EBA and NPHS cases, the solution of the field equation is of the Bogoljubov form:

$$f_k(\eta) = \alpha_k^{\text{mv}} \sqrt{-\eta} H_\nu^{(1)}(-k\eta) + \beta_k^{\text{mv}} \sqrt{-\eta} H_\nu^{(2)}(-k\eta)\,, \tag{14}$$

where $\alpha_k^{\text{mv}}, \beta_k^{\text{mv}}$ are determined by the specific boundary conditions (choice of vacuum) imposed on the solution at the time η_c, and by the Wronskian condition, which translates into $(\alpha_k^{\text{mv}})^2 - (\beta_k^{\text{mv}})^2 = 1$. In general the Bogoljubov coefficients depend on the mode k and the cutoff scale and time (these latter are related in the NPHS case, but not in the EBA, as we said).

The two-point function at small scales, at leading order in slow-roll and β_k^{mv}, is

$$P_{\text{mv}}(k) \underset{k\to 0}{\sim} \frac{H^2}{4M_{\text{Planck}}^2 \epsilon k^3}\left(1 + 2\,\text{Re}(\beta_k^{\text{mv}} e^{-i\text{Arg}(\alpha_k^{\text{mv}})})\right). \tag{15}$$

3.4 General constraints on Bogoljubov coefficients

The precise magnitude and scale (k) dependence of the coefficients $\beta_k^{\text{mdr}}, \beta_k^{\text{mv}}$, entering respectively equations (13) for the modified dispersion relations scenario, and (15) for the modified vacuum, are determined in full details by the specific model or boundary condition giving rise to the modified dispersion relation or modified vacuum. However, to be in accordance with observations and to avoid backreaction stopping the slow-roll inflationary evolution, these coefficients are subject to a series of phenomenological constraints, see for example [15, 16]:

- the observations on the spectral index show that $-k\frac{d\log(k^3P(k))}{dk}$ is small, so that $\beta_k^{\text{mdr}}, \beta_k^{\text{mv}}$ must be slowly varying with k at the observed scales;
- backreaction imposes two further constraints:
    - the total energy density must be finite. In general this demands that $|\beta_k^{\text{mdr}}|, |\beta_k^{\text{mv}}|$ decay faster than k^{-2} at large k;
    - preserving the slow-roll inflationary evolution enforces the constraint [15, 16]

$$|\beta_k^{\text{mdr}}|, |\beta_k^{\text{mv}}| \leq \sqrt{\epsilon|\mu|}\frac{HM_{\text{Planck}}}{\Lambda^2}\,, \qquad \mu \equiv \eta_{\text{sl}} - \epsilon\,. \tag{16}$$

In the following we will keep indicating the scale dependence of these coefficients with the label $_k$. observe also that we distinguish the modified vacuum scenario from that with modified dispersion relations with the labels “mv” and “mdr”.

4 Bispectrum in the squeezing limit

We will now discuss the three-point function. We start by presenting two examples of calculations for two representative interactions (with and without higher-derivatives), to show in details and step-by-step how their contributions to the bispectrum are affected by the generic features we outlined in the introduction (particle creation, presence of new scales related to new physics and/or initial time). These interactions were studied, but not in the squeezing limit, in [4, 9, 15, 16, 21].

We will then generalize our analysis to the other possible cubic interactions and see if the above mentioned generic features of the scenarios suffice to determine in general, and in what measure, the leading order form of the bispectrum. Finally, we comment on the results in section 5.

Let us introduce some notation. For the purpose of computing the bispectrum, it is convenient to use the field redefinition [4, 9, 15][4]

$$\zeta = \zeta_c + \frac{1}{8}\frac{\dot{\phi}^2}{H^2 M_{\text{Planck}}^2}\zeta_c^2 + \frac{1}{4}\frac{\dot{\phi}^2}{H^2 M_{\text{Planck}}^2}\partial^{-2}(\zeta_c \partial^2 \zeta_c) + \frac{1}{2}\frac{\ddot{\phi}}{\dot{\phi} H}\zeta_c^2 \tag{17}$$

Using the definitions of the slow-roll parameters, the bispectrum of ζ, which is the relevant one for observations, is related to the three-point function for ζ_c as [4]

$$\begin{aligned}\langle \zeta_{\vec{k}_1}(\eta)\zeta_{\vec{k}_2}(\eta)\zeta_{\vec{k}_3}(\eta)\rangle &= \langle \zeta_{c,\vec{k}_1}(\eta)\zeta_{c,\vec{k}_2}(\eta)\zeta_{c,\vec{k}_3}(\eta)\rangle \\ &\quad + (2\pi)^3\delta\Big(\sum_i \vec{k}_i\Big) \sum_{\substack{i=1 \\ \text{mod } 3}}^{3} \Big(\epsilon\Big(\frac{3}{2} + \frac{k_{i+1}^2 + k_{i+2}^2}{2k_i^2}\Big) - \eta_{\text{sl}}\Big) P(k_{i+1})\, P(k_{i+2})\,. \end{aligned} \tag{18}$$

The two-point function and the quadratic part of the action are the same for ζ and ζ_c [4].

Using the variable ζ_c, the general formula for the three-point function at leading order (tree-level) in perturbation theory is

$$\langle \zeta_c(\eta,\vec{x}_1)\zeta_c(\eta,\vec{x}_2)\zeta_c(\eta,\vec{x}_3)\rangle = -2\text{Re}\left(\int_{\eta_{\text{in}}}^{\eta} d\eta' i \langle \psi_{\text{in}}|\zeta_c(\eta,\vec{x}_1)\zeta_c(\eta,\vec{x}_2)\zeta_c(\eta,\vec{x}_3) H_{(I)}(\eta')|\psi_{\text{in}}\rangle\right)\,, \tag{19}$$

where η_{in}, $|\psi_{\text{in}}\rangle$ are the initial time and state (vacuum), and $H_{(I)}$ is the interaction Hamiltonian.

The modifications due to the scenarios we are considering will affect both the first connected and time-integrated contribution on the right-hand side of equation (18), both the other not time-integrated ones, via the modified two-point functions. However, the latter modifications, as we have reviewed in the previous sections, are strongly suppressed. We will thus focus on the time-integrated connected contribution, which is expected to be more sensitive to the modifications appearing in our scenarios, see [15, 16].

We will write the result for the three-point function in powers of $|\beta_{k_i}|$. Up to linear order,

$$\langle \zeta_{c,\vec{k}_1}\zeta_{c,\vec{k}_2}\zeta_{c,\vec{k}_3}\rangle = \delta_0\langle \zeta_{c,\vec{k}_1}\zeta_{c,\vec{k}_2}\zeta_{c,\vec{k}_3}\rangle + \delta_\beta\langle \zeta_{c,\vec{k}_1}\zeta_{c,\vec{k}_2}\zeta_{c,\vec{k}_3}\rangle \tag{20}$$

with

$$\delta_0\langle \zeta_{c,\vec{k}_1}\zeta_{c,\vec{k}_2}\zeta_{c,\vec{k}_3}\rangle = A(k_1,k_2,k_3)\delta F_0(k_1,k_2,k_3)\,, \qquad \delta_\beta\langle \zeta_{c,\vec{k}_1}\zeta_{c,\vec{k}_2}\zeta_{c,\vec{k}_3}\rangle = A(k_1,k_2,k_3)\delta F_1(k_1,k_2,k_3) \tag{21}$$

In the standard scenario we define $\delta F_0^{\text{standard}} = 1$, $\delta F_1^{\text{standard}} = 0$, so that $A(k_1,k_2,k_3)$ is indeed the standard three-point function for ζ_c. In the modified scenarios, δF_0 and δF_1 represent therefore the corrections to the standard result, respectively of order $|\beta_{k_i}|^0$ and $|\beta_{k_i}|^1$, due to the new high energy physics.

---

[4] In [15], ζ_c is simply written as ζ, see their equation (3.11). The same is done in many other papers, such as [9].

4.1 Example 1: Minimal coupling cubic interaction

We begin by studying the case of the cubic interaction [4]:

$$H_{(I)} = -\int d^3x\, a^3\, (\frac{\dot{\phi}}{H})^4 \frac{H}{M^2_{\text{Planck}}}\, \zeta_c'^2 \partial^{-2}\zeta_c' \,. \tag{22}$$

We call this interaction the "minimal coupling cubic interaction" because it is already present in the simplest case of a scalar field (inflaton) minimally coupled to gravity [4].

In this case equation (19) leads to

$$\langle \zeta_{c,\,\vec{k}_1}(\eta)\zeta_{c,\,\vec{k}_2}(\eta)\zeta_{c,\,\vec{k}_3}(\eta)\rangle = 2\mathrm{Re}\Big(-i(2\pi)^3\delta(\sum_i \vec{k}_i)\Big(\frac{\dot{\phi}}{H}\Big)^4 \frac{H}{M^2_{\text{Planck}}}\int_{\eta_{\text{in}}}^{\eta} d\eta' \frac{a(\eta')^3}{k_3^2}\prod_{i=1}^{3}\partial_{\eta'}G_{k_i}(\eta,\eta') + \text{permutations}\Big), \tag{23}$$

where $\eta \sim 0$ is a very late time when all modes k_i are outside the horizon. We will now present the results for the standard and modified scenarios. All the relevant Whightman functions are listed in appendix A.2.

4.1.1 Standard scenario (review)

In the standard scenario it is $\eta_{\text{in}} = -\infty$, and the Whightman functions are obtained from the definition (4) using the mode functions in equation (7), see appendix A.2. Inserting them in equation (23), one finds [4]

$$A(k_1,k_2,k_3) \equiv \langle \zeta_{c,\,\vec{k}_1}\zeta_{c,\vec{k}_2}\zeta_{c,\vec{k}_3}\rangle_{\text{st}} = 4(2\pi)^3\delta(\sum_i \vec{k}_i)\frac{H^6}{\dot{\phi}^2 M^2_{\text{Planck}}}\frac{k_1^2k_2^2k_3^2}{\prod_{i=1}^{3}(2k_i^3)}\sum_l \frac{1}{k_t k_l^2}\,, \qquad k_t = \sum_{i=1}^{3} k_i \,. \tag{24}$$

Adding the disconnected contribution in equation (18) to obtain the bispectrum for ζ, and taking the squeezed limit $k_1 \ll k_S$, $k_{2,3} \sim k_S$, finally

$$\langle \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\rangle_{\text{st}} \underset{k_1 \ll k_S}{\simeq} (2\pi)^3\delta(\sum_i \vec{k}_i)(1-n_s)P_{\text{st}}(k_1)P_{\text{st}}(k_S)\,. \tag{25}$$

Equation (25) is the so-called local form of the bispectrum [4–6].

4.1.2 Modified initial vacuum

As the computations are simpler in this case, we present it first. The time integration in equations (19), (23) has now a lower limit at the finite time $\eta_{\text{in}} = \eta_c$. The relevant Whightman functions are listed in appendix A.2. The result for the bispectrum for generic k_1, k_2, k_3 has been first presented in [15]. Writing the result in the form (20), (21) using (24), yields

$$\delta F_0 \sim 1\,, \qquad \delta F_1 = \sum_{j=1}^{3}\delta F_1^{(j)} = -\sum_{j=1}^{3}\frac{k_1+k_2+k_3}{\sum_{h\neq j}k_h - k_j}\mathrm{Re}\big[\beta_{k_j}^{\text{mv}*}\big(1-e^{i(\sum_{h\neq j}k_h - k_j)\eta_c}\big)\big]\,. \tag{26}$$

In the squeezing limit $k_1 \ll k_S$, $k_{2,3} \sim k_S$, the contribution proportional to β_{k_1} is very subdominant, as it is given by

$$\delta F_1^{(1)} \underset{k_1 \ll k_S}{\simeq} -\mathrm{Re}\big[\beta_{k_1}^{\text{mv}*}\big(1-e^{i2k_S\eta_c}\big)\big] \ll 1\,. \tag{27}$$

Larger contributions are found when the perturbations depending on the large momenta in the squeezing limit are in opposition of phase, that is, k_j is either k_2 or k_3. Indeed, in that case the denominator and the argument of the exponential in $\delta F_1^{(j=2,3)}$ are very small: using $k_{h\neq 1,j} = (k_1^2 + k_j^2 + 2k_1 k_j \cos\theta_j)^{\frac{1}{2}}$, from momentum conservation, and expanding in $\frac{k_1}{k_j} = \frac{k_1}{k_S}$, one obtains

$$\text{for } j = 2 \text{ or } 3 \qquad \sum_{h\neq j} k_h - k_j \underset{k_1 \ll k_{2,3}}{\simeq} k_1(1+\cos\theta_j) \equiv k_1\, v_{\theta_j} \,. \tag{28}$$

so that

$$\delta F_1^{(j=2,3)} \underset{k_1\ll k_S}{\simeq} -\frac{k_1+k_2+k_3}{k_1\, v_{\theta_j}} \mathrm{Re}\big[\beta_{k_j}^{\mathrm{mv}*}\big(1 - e^{ik_1\,\eta_c v_{\theta_j}}\big)\big] \,. \tag{29}$$

It can be easily checked (for example by plotting) that the one in (28) is an almost perfect approximation for all values of θ_j already for $\frac{k_1}{k_j} = \frac{k_1}{k_S} \sim 0.1$.

The form of the correction $\delta F_1^{(j=2,3)}$, then, depends on the interplay between the scales k_1 and η_c. Recalling that we consider the realistic limit, where k_1 is small but not zero, there are two asymptotic possibilities where the result can be evaluated most explicitly:

$$1)\ |k_1\eta_c v_{\theta_j}| \gg 1 \quad \Rightarrow \quad \delta F_1^{(j=2,3)} \underset{k_1\ll k_S}{\simeq} -2\frac{k_S}{k_1} v_{\theta_j}^{-1}\,\mathrm{Re}\big[\beta_{k_S}^{\mathrm{mv}*}\big] \tag{30}$$

$$2)\ |k_1\eta_c v_{\theta_j}| \ll 1 \quad \Rightarrow \quad \delta F_1^{(j=2,3)} \underset{k_1\ll k_S}{\simeq} -2k_S\eta_c\,\mathrm{Im}\big[\beta_{k_S}^{\mathrm{mv}*}\big] \,. \tag{31}$$

We see that these contributions dominate over (27) because of the large factors $\frac{k_S}{k_1}$ and $k_S\eta_c$. In the case 1) we have eventually taken into account that in standard observables such as the halo bias, see section 6, the final result involves an integration/average over quantities such as θ_j, k_S and so the terms with large oscillations averages to zero.

Thus, the leading correction to the squeezed limit of the three-point function for non-zero β_k reads

$$\delta_\beta\langle\zeta_{c,\vec{k}_1}\zeta_{c,\vec{k}_2}\zeta_{c,\vec{k}_3}\rangle^{\mathrm{mv}}_{\text{min cub}} = A(k_1,k_2,k_3)\,\delta F_1(k_1,k_2,k_3) \underset{k_1\ll k_S}{\simeq} (2\pi)^3\delta\big(\sum_i \vec{k}_i\big)\ \mathcal{B}^{\mathrm{mv}}_{\text{min cub}}\, P_{\mathrm{st}}(k_1)P_{\mathrm{st}}(k_S)\,, \tag{32}$$

with

$$\mathcal{B}^{\mathrm{mv}}_{\text{min cub}} = \sum_{j=2}^{3}\mathcal{B}^{\text{mv, min cub}}_{(j)}\,, \qquad \mathcal{B}^{\text{mv, min cub}}_{(j)} = \begin{cases} -4\,\epsilon\,\frac{k_S}{k_1}\,v_{\theta_j}^{-1}\mathrm{Re}\big[\beta_{k_S}^{\mathrm{mv}*}\big] & \text{if } |k_1\eta_c v_{\theta_j}| \gg 1 \\ -4\,\epsilon\,k_S\eta_c\,\mathrm{Im}\big[\beta_{k_S}^{\mathrm{mv}*}\big] & \text{if } |k_1\eta_c v_{\theta_j}| \ll 1 \end{cases} \tag{33}$$

For observational purposes, for example concerning the halo bias, one is interested in the magnitude of these corrections and in their dependence on the probed large scale k_1^{-1}. We will discuss these aspects in details in section 5. We can already notice, however, that the k_1-dependence for the leading contributions (32), (33) is fully determined, since $\beta_{k_S}^{\mathrm{mv}}$ depends on the momentum k_S and not on k_1. The magnitude of the contribution will have instead to be estimated and bound using the phenomenological constraints reviewed in section 3.4.

4.1.3 Modified dispersion relations

The bispectrum in this case was first computed in [16], to which we address the reader for more details. The Whightman functions necessary to compute the three-point function are obtained from their definition in equation (4) using the field solution (10). Hence, the time integral in the correlator (23) breaks up into the different intervals of validity of the piecewise solutions (10). In [16], it was shown that the leading contribution occurs when the Whightman functions have support in the intervals IV and III, see figure 1. Other contributions are indeed suppressed by Δ, defined in equation (11), or by a rapidly decaying integrand[5], see [16].

---

[5] The path of integration in time must be chosen such that the oscillating piece of the integrand becomes exponentially decreasing for $\eta \to \infty$. This corresponds to taking the vacuum of the interacting theory [4].

We concentrate, therefore, only on the leading contributions from intervals IV and III. It is convenient to write the result in terms of the variables

$$y \equiv -\frac{p_{\rm max}(\eta)}{\Lambda} = \frac{H}{\Lambda} k_{\rm max} \eta \,, \qquad x_i \equiv \frac{k_i}{k_{\rm max}} \,. \tag{34}$$

In particular, when taking the squeezing limit, it is $k_{\rm max} \approx k_{2,3} \approx k_S$ and $x_1 \ll 1$. Since y is a rescaled time coordinate, sometimes we will call it simply “time” in this section.

By inserting the relevant Whightman functions, listed in appendix A.2, in equation (23) and writing the result in the form (20), (21), with (24), the corrections $\delta F_{m=\{0,1\}}$ read [16]

$$\delta F_{m=\{0,1\}}(x_1, x_2, x_3, y) = \sum_{j=1}^{3} \mathrm{Re}\left[(\beta_{k_j}^{\rm mdr*})^m \frac{\Lambda}{H} \int_{y_{\rm II}}^{y} dy' x_t g(\{x_{h\neq j}\}, x_j, y', m)\, e^{i\frac{\Lambda}{H} S_0(\{x_{h\neq j}\}, x_j, y', m)} \right] , \tag{35}$$

where $x_t \equiv \sum_{i=1}^{3} x_i$, and $g(\{x_{h\neq j}\}, x_j, y', m)$ is reported in equation (106) of appendix A.3. S_0 is defined as[6]

$$S_0(\{x_{h\neq j}\}, x_j, y', m) = \int^{y'} dy'' \left(\sum_{h\neq j} \omega(x_h\,, y'') + (-1)^m \omega(x_j\,, y'') \right) , \qquad h, j \in \{1, 2, 3\} \,, \tag{36}$$

The limits of integration for the variable y' in (35) have been discussed in [16], and we refer the reader to that reference for the details, reporting here the results. The upper limit is $y \approx 0$ because the bispectrum is evaluated at late time $\eta \approx 0$, while the lower limit $y_{\rm II}$ is determined by the smallest among $\eta_{\rm II}^{(k_{i=1,2,3})}$, which bound region III for the three momenta $k_{i=1,2,3}$, see equation (10). As shown in [16], it is $y_{\rm II} = \frac{H}{\Lambda} k_{\rm max} \eta_{\rm II}^{(k_{\rm max})} \approx -1$. This is indeed straightforward: the corrections to the linear dispersion relation at the time $\eta_{\rm II}^{(k)}$ that separates region III and II must be quite important in order to drive the frequency close to the turning point (see figure 1), and so, since these corrections grow as powers of $-\frac{p(\eta)}{\Lambda}$, see (9), it must be $|p(\eta_{\rm II}^{(k)})\Lambda^{-1}| \sim 1$. Furthermore, since $p(\eta_{\rm II}^{(k)}) = -k\eta_{\rm II}^{(k)} H$, the smallest boundary time $\eta_{\rm II}^{(k)}$ is obtained for the largest k. Thus, we have to consider $\eta_{\rm II}^{(k_{\rm max})}$ and, from the definition of y, it must be $|y_{\rm II}| = |p_{\rm max}(\eta_{\rm II}^{(k_{\rm max})})\,\Lambda^{-1}| \sim 1$. Observe then that $\forall |y| \leq |y_{\rm II}|$ it is $|x_1 y| \leq |x_1 y_{\rm II}| \ll 1$ in the squeezing limit, so effectively $F(\frac{H}{\Lambda} k_1 \eta) = F(x_1 y) \simeq 1$ and $\omega(x_1 y) \simeq x_1$.

The integral is then divided in two contributions, according to whether the Whightman functions depending on $k_{2,3}$ have support in their region IV or III. In the squeezing limit, they will have support mostly in region III, because the modes will become superhorizon (region IV) only at $\eta_S \simeq -k_{2,3}^{-1} \simeq -k_{\rm max}^{-1}$, that is for $y_S = -\frac{H}{\Lambda} \sim 0$: this time is also pretty close to the late time $y \sim 0$ that the bispectrum is evaluated at, thus the contribution from when all Whightman functions have support in region IV is subdominant. We concentrate now on the contribution from region III (of course $\zeta_{\vec{k}_1}$ will have supports both in its III and IV regions).

The one in equation (35) is a typical Fourier integral, dominated by the contributions of points where the strong suppression ($\frac{\Lambda}{H} \gg 1$) from the oscillating phase of the integrand is reduced. Concerning δF_0, the only such points are the boundary ones: in fact, there are no stationary points since the first derivative of the phase $S_0(x_{1,2,3}, y', m = 0)$ is the sum of the positive-defined frequencies and thus is always positive and never zero, see equation (36) for $m = 0$. Fourier integrals like this can be approximated for $\frac{\Lambda}{H} \gg 1$ by integrating by parts (see [27] and section 4.1.2 in [16]). Thus, in the squeezing limit $x_1 \ll 1$, $x_{2,3} \simeq x_S \simeq 1$ we find

$$\delta F_0 \underset{\substack{k_1 \ll k_S \\ (x_1 \ll 1)}}{\simeq} 1 - \frac{1}{\omega(1,-1)} \mathrm{Im}\Big[g(x_1, x_2, x_3, -1, 0)\, e^{i\frac{\Lambda}{H} S_0(x_1, x_2, x_3, -1, 0)} \Big]\Big|_{\substack{x_1 \simeq 0 \\ x_{2,3} \simeq x_S \simeq 1}} \simeq 1 \,, \tag{37}$$

---

[6] Observe that $\omega(x, y) \equiv xF(xy)$ is dimensionless.

where in the last passage we have taken into account, as before, the averaging to zero of the rapid oscillations in observables such as the halo bias. As we see, this contribution does not lead to very significant corrections to the standard local form of the bispectrum.

We expect a more interesting behaviour for what concerns δF_1 [16]. In the squeezing limit $x_1 \ll 1$, we find that the dominant contributions are those where the mode function in the negative-energy branch is the one depending on k_2 or k_3, because in those cases the suppression from the oscillations is very reduced, since the perturbations depending on the large momenta are in opposition of phase:

for $j = 2$ or 3:

$$
\begin{aligned}
S_0(\{x_{h\neq j}\}, x_j, y', 1) &= \int^{y'} dy'' \Big(\sum_{h\neq j} \omega(x_h\,, y'') - \omega(x_j\,, y'') \Big) \underset{x_1 \ll 1}{\simeq} x_1 \int^{y'} dy'' \big(1 + \partial_x \omega(x, y'')\big|_{x_S} \cos\theta_j\big) + \mathcal{O}(x_1^2) \\
&= x_1\, y'(1 + F(y'))\cos(\theta_j) + \mathcal{O}(x_1^2) \quad \equiv \; x_1\, \tilde{v}_{\theta_j}(y')\,, \qquad (38)
\end{aligned}
$$

where we have used equation (9) and $x_s \simeq 1$.

The solution for the integral has two asymptotic regimes, depending on the magnitude of the phase factor $\frac{\Lambda}{H}\, x_1$ and the behaviour of the integrand at boundary and stationary points, see [27]. We have listed in appendix A.3, table 1, the relevant information, which leads to the solution

$$
\delta F_1 \simeq
\begin{cases}
\left\{
\begin{array}{l}
\left[\begin{array}{ll}
-\frac{2}{x_1}\sum\limits_{j=2}^{3} \frac{\mathrm{Re}\left[\beta^{\mathrm{mdr}^*}_{k_S}\right]}{1+\cos(\theta_j)} & \text{if } v_{\theta_j} > \frac{H}{\Lambda}\frac{1}{x_1} \\
\left(\frac{\Lambda}{H}\right)^{1\text{-}\frac{1}{\kappa+1}} \frac{4\Gamma(\frac{1}{\kappa+1})}{x_1^{\frac{1}{\kappa+1}}} \frac{\mathrm{Im}\left[\beta^{\mathrm{mdr}^*}_{k_S}\, e^{i\frac{\pi}{2}\frac{\mathrm{sign}(F^{(\kappa)})}{\kappa+1}}\right]}{(\kappa+1)|F^{(\kappa)}|^{\frac{1}{\kappa+1}}} & \text{if } v_{\theta_j} < \frac{H}{\Lambda}\frac{1}{x_1}
\end{array}\right]
- \frac{2}{x_1}\sum\limits_{j=2}^{3} \frac{\mathrm{Im}\left[\beta^{\mathrm{mdr}^*}_{k_S}\, g(\text{-}1) e^{i\frac{\Lambda}{H}x_1 \tilde{v}_{\theta_j}(\text{-}1)}\right]}{\tilde{v}^{(1)}_{\theta_j}(\text{-}1)} \\
\qquad + \sum\limits_{j=2}^{3} \sum\limits_{\{y_*\}} \left(\frac{\Lambda}{H}\right)^{1\text{-}\frac{\mu_*}{\nu_*}} \frac{1}{x_1^{\frac{\mu_*}{\nu_*}}} \mathrm{Re}\left[\beta^{\mathrm{mdr}^*}_{k_S}\, \mathcal{I}^{(\mu_*,\nu_*)}_{\theta_j}(y_*)\, e^{i\frac{\Lambda}{H}x_1 \tilde{v}_{\theta_j}(y_*)}\right]
\end{array}
\right\} & \text{if } \frac{\Lambda}{H}x_1 \gg 1 \\[3em]
2\frac{\Lambda}{H}\, \mathrm{Im} \sum\limits_{j=2}^{3} \left[\beta^{\mathrm{mdr}^*}_{k_S} (\mathcal{O}(1) + \mathcal{O}(x_1))\right] & \text{if } \frac{\Lambda}{H}x_1 \ll 1
\end{cases}
$$

This complete leading-asymptotic formula looks complicated, and we write it only this time for illustrative purposes, but the final result will be simpler and clearer. The notation is explained in appendix A.3. Let us recall here the main points. We have defined

$$
\mathcal{I}^{(\mu_*,\nu_*)}_{\theta_j}(y_*) = \sum_{j=2}^{3} \frac{\Gamma(\frac{\mu_*}{\nu_*})}{\nu_* \big(|\tilde{v}^{(\nu_*)}_{\theta_j}(y_*)|\big)^{\frac{\mu_*}{\nu_*}}}\, g^{(\mu_*\text{-}1)}(y_*) \Big(\sum_{l=0}^{1} (\text{-}1)^{l(\mu_*\text{-}1)} e^{(\pm 1)^l i \frac{\pi}{2}\frac{\mu_*}{\nu_*}\mathrm{sign}(\tilde{v}^{(\nu_*)}_{\theta_j}(y_*))}\Big) \qquad (39)
$$

where, $\tilde{v}^{(\nu_*)}_{\theta_j}(y_*)$, $g^{(\mu_*\text{-}1)}(y_*)$ are the coefficient of the leading behaviour of the functions $\tilde{v}_{\theta_j}(y)$, $g(\{x\}, y)$ in the squeezing limit in proximity of a stationary point y_*, see table 1. The sign ± in the phases depends on $\tilde{v}_{\theta_j}(y) - \tilde{v}_{\theta_j}(y_*)$ being even or odd under $(y - y_*) \to -(y - y_*)$, for $y \sim y_*$. If ν_* is an integer, $\tilde{v}^{(n)}_{\theta_j} = n!^{-1}\, \partial^n_{y'} \tilde{v}_{\theta_j}$ (similarly for μ_*). Recall also that $v_{\theta_j} \equiv 1+\cos(\theta_j)$.

Finally, the contribution of the (nearly) folded configurations (singled out by the condition $v_{\theta_j} \equiv 1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{1}{x_1}$) is determined by the lowest order correction to the dispersion relation, which, expanding (9), would go (with model-dependent $F^{(\kappa)} \sim \mathcal{O}(1)$ and κ) as

$$
\omega_{\text{phys}}(p) \sim p\big(1 + \big(\frac{p}{\Lambda}\big)^{\kappa} F^{(\kappa)} + \cdots\big)\,. \qquad (40)
$$

In equation (39), the contributions in the first line for $\frac{\Lambda}{H}x_1 \gg 1$ arise from the boundary points, the second line from the possible presence of a number of stationary points $\{y_*\}$, which for generality we have taken as stationary of order ν_* for $\tilde{v}_{\theta_j}(y)$, and as zeroes of the function g of order μ_*-1, see [27] (μ_* and ν_* are generally different for the various points). The presence of stationary points is strictly dependent on the specific dispersion relations, that is the specific model. However, their contributions will be suppressed in the final observables such as the halo bias, because of the averaging to zero of their large oscillations, visible in (39).

Inserting these results in equation (20), and using (21), (24), the leading correction to the standard consistency relation due to non-zero β_k reads

$$\delta_\beta \langle \zeta_{c,\vec{k}_1} \zeta_{c,\vec{k}_2} \zeta_{c,\vec{k}_3} \rangle^{\text{mdr}}_{\text{min cub}} \underset{k_1 \ll k_S}{\simeq} (2\pi)^3 \delta(\sum_i \vec{k}_i)\, \mathcal{B}^{\text{mdr}}_{\text{min cub}}\, P_{\text{st}}(k_1) P_{\text{st}}(k_S)\,, \tag{41}$$

$$\mathcal{B}^{\text{mdr}}_{\text{min cub}} = \sum_{j=2}^{3} \begin{cases} \begin{bmatrix} -4\,\epsilon\,\frac{k_S}{k_1}\,\frac{1}{1+\cos(\theta_j)}\text{Re}\left[\beta^{\text{mdr}*}_{k_S}\right] & \text{if } 1+\cos(\theta_j) > \frac{H}{\Lambda}\frac{1}{x_1} \\ 4\,\epsilon\left(\frac{\Lambda}{H}\right)^{1-\frac{1}{\kappa+1}} \left(\frac{k_S}{k_1}\right)^{\frac{1}{\kappa+1}} \Gamma\left(\frac{1}{\kappa+1}\right) \frac{\text{Im}\left[\beta^{\text{mdr}*}_{k_S} e^{i\frac{\pi}{2}\frac{\text{sign}(F^{(\kappa)})}{\kappa+1}}\right]}{(\kappa+1)|F^{(\kappa)}|^{\frac{1}{\kappa+1}}} & \text{if } 1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{1}{x_1} \end{bmatrix} & \text{if } \frac{\Lambda}{H}x_1 \gg 1 \\ \\ 4\,\epsilon\,\frac{\Lambda}{H}\,\text{Im}\left[\beta^{\text{mdr}*}_{k_S}\,\mathcal{O}(1)\right] & \text{if } \frac{\Lambda}{H}x_1 \ll 1 \end{cases} \tag{42}$$

where we have accounted for the averaging to zero of the rapid oscillations in (39) in observables such as the halo bias. Again, at leading order the dependence on k_1 of the result is fully specified, and there appear enhancement factors such as $\frac{\Lambda}{H}$ and $\frac{k_S}{k_1}$. We will comment on the result in section 5.

The final result is very similar to the one obtained in the modified vacuum case, which points at the fact that indeed the squeezing limit is dominated, at leading order, by the generic features common to both scenarios (such as particle creation, interference effects). The only difference is that in the modified vacuum case, the folded configurations yield a contribution corresponding to $\kappa \to \infty$ in (42), since the phase of the integrand and all its derivatives are identically zero, see (33). In the case of modified dispersion relations, instead, the true analogous of the folded configuration (which would be $\sum_{h \neq j} \omega(x_h\,, y') - \omega(x_j\,, y') = 0$ for some x's but *for all* y'*'s*) does not occur[7] and κ is finite.

4.2 Example 2: Quartic derivative interaction

In the standard scenario, the contributions to the bispectrum from higher derivative interactions are suppressed in the squeezing limit by powers of $\frac{k_1}{k_S}$ [6]. It is interesting, then, to study if the additional contributions to non-Gaussianities due to particle creation in the scenarios of modified vacuum and modified dispersion relations would lead to the same suppressed contributions to the bispectrum and the same overall scale dependence.

In this section, we present a detailed example of the kind of calculations involved, before passing to the more general discussion in section 4.3. We take as our representative example the cubic interaction coming from the correction $\mathcal{L}_{\text{HDI}} = \sqrt{-\text{det}G}\frac{g}{8M^4}((\nabla\Phi)^2)^2$ to the effective action of the inflaton Φ, where G is the metric, g the coupling. In the standard single-field slow-roll scenario, [21] has discussed that this is the lowest-dimensional operator in the class of operators which is the most important for non-Gaussianities. Nonetheless, for the standard scenario its contribution is very suppressed in the squeezing limit, as we have said.

---

[7] It can be easily verified at least if ω is given by elementary functions and/or power series.

Expanding in perturbations and converting in terms of ζ as explained in details in [15, 16], from $\mathcal{L}_{\text{HDI}}$ one obtains the cubic term

$$H_{(I)} = -\int d^3x a \frac{g\,\dot{\phi}^4}{2H^3\Lambda^4}\zeta_c'\big(\zeta_c'^2 - (\partial_i\zeta_c)^2\big)\,. \tag{43}$$

Because of its origins from the term $\mathcal{L}_{\text{HDI}}$, which contains quartic power of derivatives, we call this cubic interaction "quartic derivative interaction". It is allowed to treat these interactions perturbatively also when the physical momenta $p = a^{-1}k$ are such that $\frac{p}{\Lambda}$ is not small any more, because, compared to the operators changing the dispersion relation, they are further suppressed by additional powers of the perturbation fields.

The bispectrum, from equation (19), reads in this case

$$\begin{aligned}\langle\zeta_{c,\vec{k}_1}(\eta)\zeta_{c,\vec{k}_2}(\eta)\zeta_{c,\vec{k}_3}(\eta)\rangle = 2\text{Re}\Bigg[&-i(2\pi)^3\delta^{(3)}\big(\sum_i \vec{k}_i\big)\frac{g\dot{\phi}^4}{2H^3\Lambda^4}\int_{\eta_{in}}^{\eta} d\eta'\, a\big(\partial_\eta G_{k_1}(\eta,\eta')\partial_\eta G_{k_2}(\eta,\eta')\partial_\eta G_{k_3}(\eta,\eta') + \\ &+ \vec{k}_1\cdot\vec{k}_2 G_{k_1}(\eta,\eta')G_{k_2}(\eta,\eta')\partial_\eta G_{k_3}(\eta,\eta')\big) + \text{permutations}\Bigg]\,,\end{aligned} \tag{44}$$

where, again, $\eta \sim 0$ is a very late time when all modes k_i are outside the horizon.

4.2.1 Modified initial state

In this scenario the leading contribution to the bispectrum for generic external momenta has been computed in [9, 15]. Writing once more the result in the form (20), (21), with (24), it is found

$$\delta F_1(k_1,k_2,k_3,\eta\simeq 0) = \text{Re}\left[iC\frac{k_S^3}{k_1^2k_2^2k_3^2}\sum_{j=1}^{3}\beta_{k_j}^{\text{mv}*}\int_{\eta_c}^{0} d\eta' e^{i(\sum_{h\neq j}k_h - k_j)\eta'}P_j(k_1,k_2,k_3,\eta') + \text{c.c.}\right], \tag{45}$$

with

$$C \equiv \frac{x_t\prod_{l=1}^{3}x_l^2}{\sum_{r>v}x_r^2x_v^2}\frac{gH^2M_{\text{Planck}}^2}{4\Lambda^4}\,. \tag{46}$$

$$\begin{aligned}P_j(k_1,k_2,k_3,\eta') = -k_t\prod_{i=1}^{3}\big(\sum_{h\neq i}k_h - k_i\big) - i\eta'\Big(&(k_{j+1}^2-k_j^2-k_{j+2}^2)k_{j+1}^2(k_{j+2}-k_j) + (k_{j+2}^2-k_j^2-k_{j+1}^2) \\ &k_{j+2}^2(k_{j+1}-k_j) + (k_j^2-k_{j+1}^2-k_{j+2}^2)k_j^2(k_{j+1}+k_{j+2})\Big) + (-\eta')^2\big(\sum_{h\neq j}k_h - k_j\big)\big(\prod_{i=1}^{3}k_i\big)(k_t^2 - 4k_{j+1}k_{j+2})\,,\end{aligned} \tag{47}$$

where j is defined modulo 3. The factor C has been written in terms of the variables x_i defined in equation (34).

As in section 4.1.2, we find that in the squeezing limit the dominating contributions to the integral occur for $j = 2, 3$ in the sum in equation (45), in particular, the largest ones come from the terms of order η and η^2 in P_j. Their integration yields

$$\int_{\eta_c}^{0} d\eta' e^{i(\sum_{h\neq j}k_h - k_j)\eta'}\eta' = \frac{1 + e^{ik_1\eta_c v_{\theta_j}}(-1 + ik_1\eta_c v_{\theta_j})}{k_1^2 v_{\theta_j}^2} \equiv \frac{I_1^{(j)}}{k_1^2 v_{\theta_j}^2}\,, \tag{48}$$

$$\int_{\eta_c}^{0} d\eta' e^{i(\sum_{h\neq j}k_h - k_j)\eta'}\eta'^2 = i\frac{2 + e^{ik_1\eta_c v_{\theta_j}}(-2 + 2ik_1\eta_c v_{\theta_j} + k_1^2\eta_c^2 v_{\theta_j}^2)}{k_1^3 v_{\theta_j}^3} \equiv \frac{I_2^{(j)}}{k_1^3 v_{\theta_j}^3}\,, \tag{49}$$

so that

$$\delta F_1 \underset{k_1 \ll k_S}{\simeq} \sum_{j=2}^{3} \frac{2gH^2 M^2_{\text{Planck}}}{\Lambda^4} \frac{k_S}{k_1} v_{\theta_j}^{-2} \left(-\text{Im}\big[\, \beta_{k_j}^{\text{mv}*}\, I_2^{(j)} \,\big] - \cos(\theta_j)\text{Re}\big[\, \beta_{k_j}^{\text{mv}*}\, I_1^{(j)} \,\big] \right) \tag{50}$$

Thus, looking at (48), (49), we find again two asymptotic behaviours for the correction to the squeezed limit of the three-point function depending on whether $|k_1 \eta_c v_{\theta_j}| \gg 1$ or $|k_1 \eta_c v_{\theta_j}| \ll 1$:

$$\delta_\beta \langle \zeta_{c,\vec{k}_1} \zeta_{c,\vec{k}_2} \zeta_{c,\vec{k}_3} \rangle^{\text{mv}}_{\text{quart der}} \underset{k_1 \ll k_S}{\simeq} (2\pi)^3 \delta(\sum_i \vec{k}_i)\ \mathcal{B}^{\text{mv}}_{\text{quart der}}\, P_{\text{st}}(k_1) P_{\text{st}}(k_S)\,, \tag{51}$$

with

$$\mathcal{B}^{\text{mv}}_{\text{quart der}} = \begin{cases} -4\,\epsilon\, \frac{gH^2 M^2_{\text{Planck}}}{\Lambda^4} \frac{k_S}{k_1} \sum\limits_{j=2}^{3} v_{\theta_j}^{-2} (2+\cos(\theta_j)) \text{Re}\big[\, \beta_{k_S}^{\text{mv}*} \,\big] & \text{if } |k_1 \eta_c v_{\theta_j}| \gg 1 \\ 2\,\epsilon\, \frac{gH^2 M^2_{\text{Planck}}}{\Lambda^4} \frac{k_1}{k_S} (k_S \eta_c)^2 \sum\limits_{j=2}^{3} \Big(\frac{2}{3} \frac{k_1}{k_S} (k_S \eta_c) v_{\theta_j}\, \text{Im}\big[\, \beta_{k_S}^{\text{mv}} \,\big] + \cos(\theta_j)\, \text{Re}\big[\, \beta_{k_S}^{\text{mv}} \,\big] \Big) & \text{if } |k_1 \eta_c v_{\theta_j}| \ll 1\,. \end{cases} \tag{52}$$

In the first of (52), we have again taken into account the averaging of large oscillations to zero in observables such as the halo bias. Recall that $|k_S \eta_c| \gg 1$ (in the NPHS scenario, in particular, $|k_S \eta_c| = \frac{\Lambda}{H}$). We comment on the actual magnitude and scale dependence of these corrections to the bispectrum in section 5.

4.2.2 Modified dispersion relations

The three-point scalar correlator for generic k_1, k_2, k_3 in this scenario was computed in [16]. We focus on the most dominant contribution, which occurs when the Whightman functions have support in regions IV/III, see (10), as discussed in [16] and reviewed in section 4.1.3. Using in equation (44) the Whightman functions listed in appendix A.2, and writing the result in the form (20), (21) with the variables in (34), the leading correction linear in $|\beta_k^{\text{mdr}*}|$ is again given by a Fourier integral, see [16]:

$$\delta F_1 = -\sum_j \text{Re}\left[C \left(\frac{\Lambda}{H} \right)^3 \beta_{k_j}^{\text{mdr}*} \int_{y_{\text{II}}}^{y} dy' y'^2\, e^{i \frac{\Lambda}{H} S_0(\{x_h\}, x_j, y', 1)}\, q(\{x_h\}, x_j, y') \right] \qquad h \neq j\,, \tag{53}$$

where the limits of integration are those presented in section 4.1.3. The function q is reported in equation (109) of appendix A.3, while S_0 and C are defined respectively in equation (36) and (46).

When taking the squeezing limit, the dominant contributions are again those where the perturbations depending on k_2, k_3 are in opposition of phase. Then, $S_0(\{x_{1,2(3)}\}, x_{3(2)}, y', 1)$ behaves as in equation (38). The integral can be solved by asymptotic techniques, and we obtain at leading order

$$\delta_\beta \langle \zeta_{c,\vec{k}_1} \zeta_{c,\vec{k}_2} \zeta_{c,\vec{k}_3} \rangle^{\text{mdr}}_{\text{quart der}} \underset{k_1 \ll k_S}{\simeq} (2\pi)^3 \delta(\sum_i \vec{k}_i) \mathcal{B}^{\text{mdr}}_{\text{quart der}}\, P_{\text{st}}(k_1) P_{\text{st}}(k_S)$$

with (recalling appendix A.3 and table 1)

$$\begin{aligned} \mathcal{B}^{\text{mdr}}_{\text{quart der}} = -4\epsilon \frac{gH^2 M^2_{\text{Planck}}}{\Lambda^4} & \\ \times \sum_{j=2}^{3} & \begin{cases} \begin{bmatrix} \frac{k_S}{k_1} \frac{\Gamma(3)+\cos(\theta_j)}{(1+\cos(\theta_j))^2} \text{Re}\big[\, \beta_{k_S}^{\text{mdr}*} \big] & \text{if } v_{\theta_j} > \frac{H}{\Lambda} \frac{1}{x_1} \\ \frac{\Lambda}{H} \Big(\frac{\Lambda}{H} \frac{k_1}{k_S} \Big)^{1-\frac{2}{\kappa+1}} \frac{\Gamma(\frac{\kappa+3}{\kappa+1})}{(\kappa+1)} \frac{2\,\text{Im}\Big[\, \beta_{k_S}^{\text{mdr}*} F^{(\kappa)}\, e^{i \frac{\pi(\kappa+3)}{2(\kappa+1)} \text{sign}(F^{(\kappa)})} \Big]}{|F^{(\kappa)}|^{\frac{\kappa+3}{\kappa+1}}} & \text{if } v_{\theta_j} < \frac{H}{\Lambda} \frac{1}{x_1} \end{bmatrix} & \text{if } \frac{\Lambda}{H} x_1 \gg 1 \\ \frac{k_1}{k_S} \big(\frac{\Lambda}{H} \big)^2 \Big(-\frac{k_1}{k_S} \frac{\Lambda}{H} \text{Re}\big[\, \beta_{k_S}^{\text{mdr}*}\, \mathcal{O}(1) \big] + \cos(\theta_j) \text{Re}\big[\beta_{k_S}^{\text{mdr}*} \big] \Big) & \text{if } \frac{\Lambda}{H} x_1 \ll 1 \end{cases} \end{aligned} \tag{54}$$

where, again, we have taken into account the averaging to zero of the strongly oscillating terms in the final observables as the halo bias. Equation (54) matches (52); in particular, note that in the modified vacuum scenario, as we said, the folded configuration has $\kappa \to \infty$, and the dominating term in the second line of (52) is the one proportional to $\mathrm{Re}[\beta^{\mathrm{mv}}_{k_S}]$, since $|k_1\eta_c v_{\theta_j}| \ll 1$.

4.3 General analysis

Let us first summarize the key-points concerning the violation of the standard result for non-Gaussianities in the squeezing limit, which have emerged from the detailed examples discussed above:

- non-Gaussianities can be enhanced[8] in the limit by two effects:
  - interference, which reduces the suppression due to the oscillating phase of the integrand in the time integral in the formula of bispectrum, and is strongest when the perturbations depending on the large modes $k_{2,3} \sim k_S$ are in opposition of phase,
  - accumulation in time, which leads to larger enhancement from the time integration when interactions scale with higher powers of $\frac{1}{a(\eta')} \sim \eta'$ [9].
- if the probed squeezed mode and the time/scale of new physics are such that $|k_1\eta_c| \ll 1$, or $\frac{k_1}{k_S}\frac{\Lambda}{H} \ll 1$, then the perturbation depending on k_1 is outside the horizon already at the time/scale of the new physics, and thus the result is still of the local form but with enhanced non-Gaussianities (because of the interference and particle production for the other two modes $k_{2,3}$)
- if instead $|k_1\eta_c| \gg 1$ or $\frac{k_1}{k_S}\frac{\Lambda}{H} \gg 1$ (in a realistic squeezing limit $k_1 \neq 0$), the perturbation depending on k_1 is inside the horizon at the time/scale of the new physics and the bispectrum is *not* of the local form

We want now to generalize our analysis beyond the two examples of interactions discussed above. It is convenient to adopt a different gauge than the one we have used so far: in the new gauge it is $\zeta = 0$, and the perturbations are accounted for by a "matter field" perturbation φ. This gauge is fully equivalent, of course, to the one used before, and it is useful because we will have to consider only the matter action when expanding in perturbations. In particular, we can choose φ to be the inflaton perturbation, expanding the inflaton in background plus perturbations[10]

$$\Phi(x,t) = \phi(t) + \varphi(x,t)\,. \tag{55}$$

The results in terms of the variable ζ are then obtained at the end by performing a gauge transformation as described in [4]:

$$\zeta = -H\frac{\varphi}{\dot\phi} + \mathcal{O}\big(\epsilon^2, \varphi^2\big). \tag{56}$$

so that

$$\langle\zeta\zeta\zeta\rangle = -\frac{H^3}{\dot\phi^3}\langle\varphi\varphi\varphi\rangle + \Big[\langle\mathcal{O}\big(\epsilon^2,\varphi^2\big)\varphi\varphi\rangle + \text{permutations}\Big]. \tag{57}$$

---

[8] The actual magnitude of the enhancements depends on the precise value of $|\beta_{k_S}|$ and the scales of "new" physics, and thus can only be estimated, as we will show, in our phenomenological approach.

[9] This is quite intuitive, since the contribution we are interested in will arise at early times (large $|\eta'|$) around η_c for the modified vacuum scenario, or around the time of WKB violation/particle creation $\eta_{\mathrm{II}} \sim \Lambda k_S^{-1} H^{-1}$ for the modified dispersion scenario. More specifically, in the cases where $|k_1\eta_c| \ll 1$ or $\frac{k_1}{k_S}\frac{\Lambda}{H} \ll 1$, since the integral for the bispectrum becomes approximately of the form $\int_{\eta_{c,\mathrm{II}}}^0 d\eta'\,\eta'^n \sim \eta_{c,\mathrm{II}}^n$, one can easily see the growth with n. We will soon show a similar growth of the enhancement also in the cases where $|k_1\eta_c| \gg 1$, $\frac{k_1}{k_S}\frac{\Lambda}{H} \gg 1$.

[10] This gauge also makes clear from the outset the actual order in slow-roll parameters of the cubic couplings, see [4].

In particular, it is enough to consider the gauge-transformation at leading linear order, because the higher-order corrections will not yield time-integrated contributions to the bispectrum at tree-level, and, as it appears from the previous sections, it is the time-integrated contribution that leads to the leading modified results (see also [9, 15, 16]).

The most generic effective action for the inflaton Φ in single-field slow-roll inflation has been presented in [28] [11]. In a Lorentz invariant theory, the terms in the action will be function of the Lorentz invariant objects Φ, $g^{\mu\nu}\partial_\mu\Phi\partial_\nu\Phi$, $R^{\mu\nu}\partial_\mu\Phi\partial_\nu\Phi$, $\Box\Phi$, R. In particular, those that are functions of Φ only, and not of the objects involving derivative, constitute the scalar potential $V(\Phi)$. Recall also that all second order time derivatives (and first order derivatives of auxiliary fields) in higher order terms in the action (more than quadratic in the field) should be eliminated using the equation of motion, see [28]. This applies for example to $\Box\Phi$ and $R^{\mu\nu}$. In a Lorentz-broken theory (such as it is the case with modified dispersion relations), there are also additional terms that are not built up only from the Lorentz-invariant operators listed above. These terms involve higher derivatives of the fields that can be written as purely spatial derivatives in a convenient system of coordinates [12].

By expanding the effective action in the inflaton perturbation (55), given the fundamental building blocks listed above (including the higher-derivative Lorentz-breaking ones), and using conformal time, the cubic couplings for the perturbation φ have the most general and schematic form

$$\int d\eta d^3x\, a^4\, \frac{\lambda_{nms}}{\Lambda^{n+m+s-1}}\frac{\partial^n\varphi}{a^n}\frac{\partial^m\varphi}{a^m}\frac{\partial^s\varphi}{a^s} \tag{58}$$

where ∂ can be either a spatial or time derivative (comoving coordinates) [13], recalling however that in the case of time derivatives, their order is at most 1 (as we said higher orders must be eliminated by using the equations of motions, see [28]). We have sloppily indicated both time and space derivatives with the same symbol here for notational simplicity and because it will simplify the counting of powers of momenta in the correlators, but we will distinguish more carefully the two kinds of derivatives in the calculations. Finally, the dimensionless coefficients λ_{nms} can depend on $\dot{\phi}$ and in general on the slow-roll parameters, and are constrained by backreaction.

Let us comment on the possible values of n, m, s in the cubic couplings. In a Lorentz-invariant theory, given the Lorentz-invariant objects listed above, n, m, s can be at most equal to 1 [14], while in a Lorentz-broken theory their values can be higher (for the higher spatial derivatives couplings, in the convenient frame). Observe that in a gauge-fixed theory, when solving for the gravitational constraints (for example those related to the lapse N and shift-vector N^i in the ADM formalism) there will also appear non-local operators. In the gauge we are using, they arise because of the solution[15] $N^i \simeq \epsilon^{\frac{3}{2}}\partial^i(\vec{\partial})^{-2}\dot{\varphi} + \mathcal{O}(\epsilon^2)$, see [4]. This means that, when expanding in perturbations, n, m, s can also assume value -1, because of the combination $(\vec{\partial})^{-2}\dot{\varphi}$, in the leading order contribution from N^i, relevant for the bispectrum. However this combination is acted upon by ∂^i, which compensates the -1 with the additional derivative.

Observe also that N^i enters the action only via the metric $g^{\mu\nu}$ or the extrinsic curvature. In the first case, the spatial index of N^i will result always contracted with a spatial derivative, as it can be seen from $g^{\mu\nu}\partial_\mu\Phi\partial_\nu\Phi$ considering that the shift vector enters the components $g^{0i} = N^{-2}N^i$ and $g^{ij} = N^{-2}N^iN^j$. Also when coming from the extrinsic curvature, a component N^i will always be accompanied by a spatial derivative, because the extrinsic curvature is defined as $K_{ij} = (2N)^{-1}(\dot{h}_{ij} - 2\,{}^{(3)}\nabla_{\{i}N_{j\}}) = (2N)^{-1}(\dot{h}_{ij} - h_{ik}\partial_j N^k - h_{jk}\partial_i N^k - N^k\partial_k h_{ij})$, where h_{ij}, ${}^{(3)}\nabla$ are the ADM three-metric and relative covariant derivative. These properties imply that in a coupling where originally N^i was present, the combination $\partial^i(\vec{\partial})^{-2}\dot{\varphi}$ will multiply at least one other factor with a spatial derivative.

---

[11] Another way of writing the most generic effective action for a "matter scalar" takes advantage of the so-called Stueckelberg trick using the Goldstone mode for the broken time diffeomorphisms in the unitary gauge, see [29].

[12] Lorentz-breaking always implies a privileged frame, the convenient system of coordinates is precisely the one adapted to this frame.

[13] Recall that, since we are considering only scalar perturbations, after the expansion spatial derivatives indexes must be contracted by the metric, leading thus to contributions proportional, by permutation, to $\vec{k}_1 \cdot \vec{k}_2$, $\vec{k}_1 \cdot \vec{k}_3$, $\vec{k}_2 \cdot \vec{k}_3$. In particular, in the squeezing limit the contributions proportional to $\vec{k}_1 \cdot \vec{k}_2$ and $\vec{k}_1 \cdot \vec{k}_3$ give an overall contribution $\vec{k}_1 \cdot (\vec{k}_2 + \vec{k}_3) \sim k_1^2$ [6], which is subdominant with respect to the contribution $\vec{k}_2 \cdot \vec{k}_3$ that goes as $\vec{k}_2 \cdot \vec{k}_3 \to -k_2 k_3 = -k_S^2$ in the limit. The latter is thus the one we consider as leading in the following.

[14] Obviously, integrating by parts, derivatives can be "moved around", in which case the more appropriate condition is that $\max(n + m + s) = 3$. What we will say is valid, of course, also for the terms obtained by integration by parts.

[15] The standard notation $(\vec{\partial})^{-2}$ for the inverse of $\sum_j \partial^j\partial_j$ is formal. It becomes clear after a Fourier transform.

Furthermore, couplings with $n = m = s = 0$ arise from the expansion of terms such as

$$V(\Phi) \to \frac{V'''(\phi)}{3!}\varphi^3\,, \qquad \text{or} \qquad \left(\frac{\partial^r\Phi}{a^r}\right)^u \Phi^m \to (\partial_t^r\phi)^u\phi^{m-3}\varphi^3\,, \tag{59}$$

and thus are higher-order in slow-roll and would give a negligible contribution to the bispectrum (see also [4, 28, 29]). We will neglect them in the following. Moreover, couplings of the form $\frac{\lambda_{011}}{\Lambda}\varphi(\vec{\partial}\varphi)^2$, where $\vec{\partial}$ denotes spatial derivatives, are reabsorbed (eliminated) by field redefinitions such as $\phi \to \phi + \frac{\lambda_{011}}{2\Lambda}\phi^2$. Finally, couplings with a single time or spatial derivative are absent because of isotropy.

The two detailed examples that we have analyzed previously fall into the scheme we are describing. In particular, the "quartic derivative" coupling (43) derives precisely from the inflaton action, as discussed in section 4.2 and is actually the sum of two elementary cubic couplings (one with only time derivatives, the other with one time and two space derivatives). In the "minimal cubic" coupling (22), instead, we find the combination $(\vec{\partial})^{-2}\partial_\eta$ that we have been discussing. Summarizing, once written in terms of the inflaton perturbation, those couplings are of the type

$$\text{minimal coupling :} \qquad \{n, m, s\} = \{1, 1, -1\}, \qquad \lambda^{\text{min cub}}_{\text{11-1}} = \sqrt{2\epsilon}\frac{H}{M_{\text{Planck}}} \tag{60}$$

$$\text{quartic derivative :} \qquad n = m = s = 1, \qquad \lambda^{\text{quart der}}_{111} = \sqrt{\frac{\epsilon}{2}}\,g\frac{HM_{\text{Planck}}}{\Lambda^2}\,. \tag{61}$$

Finally, not writing the momenta conserving delta function, 2π factors and overall minus signs, the contribution to the bispectrum from equation (58) for a generic cubic coupling reads

$$\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\rangle_\lambda \propto \frac{\dot{\phi}^3}{H^3}\frac{\lambda_{nms}}{\Lambda^{n+m+s-1}}\int_{\eta_i}^0 d\eta'\; a(\eta')^{4-n-m-s}\;(\partial^n G)_{k_1}(\partial^m G)_{k_2}(\partial^s G)_{k_3} + \text{permutations}$$
$$+\,\text{complex conjugate} \tag{62}$$

where we have used equation (56) to cast the result in terms of the perturbation ζ (the Whightman function in this formula are those of the ζ-variable). $(\partial^r G)_k$ indicates the Fourier transform of $\partial^r G$.

We now study the dominant contributions in the different modified scenarios, recalling that, as it has emerged from the detailed examples, it occurs when the interference effects are stronger –that is, when the perturbations depending on the large modes $k_{2,3} \sim k_S$ are in opposition of phase–, and for the terms in the integrand with higher powers of η' (higher powers of $\frac{1}{a} \sim \eta'$) and, obviously, lower powers of k_1. We will not pay attention to numerical factors of order $\mathcal{O}(1)$ in the formulas as they are irrelevant for our considerations.

4.3.1 Modified vacuum, Lorentz unbroken

The relevant Whightman functions go as, see appendix A.2,

$$(\partial_{\eta'}G^{\pm})_k(0,\eta') = \pm\frac{H^4}{\dot{\phi}^2}k^2\eta'\frac{e^{\pm ik\eta'}}{2k^3}, \quad (\partial^{r\geq 0}_{x^i}G^{\pm})_k(0,\eta') = \pm\frac{H^4}{\dot{\phi}^2}(\pm ik_i)^r(1 \mp ik\eta')\frac{e^{\pm ik\eta'}}{2k^3}, \tag{63}$$

so that the leading contribution to (62), according to our analysis, is

$$\delta_\beta\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\rangle_\lambda^{\rm mv}\propto \frac{\lambda_{nms}}{\Lambda^{n+m+s\text{-}1}}\sum_{j=2}^{3}\beta^*_{k_j}\frac{H^{n+m+s+5}}{\dot\phi^3 2^3 k_1^3k_2^3k_3^3}k_a^{n+1}k_b^{m+1}k_c^{s+1}\int_{\eta_c}^{\simeq 0}d\eta'(\text{-}\eta')^{\text{-}1+n+m+s}\Big(1+(\text{-}ik_1\eta')^{n_t^{(1)}\text{-}1}\Big) \times\frac{e^{ik_1(1+\cos\theta_j)\eta'}}{2^{n_t^{(1)}}}+\text{hermitian},\quad (64)$$

where we have used equation (28) and a, b, c run over $1, 2, 3$ accounting for the permutations, from which we take the dominant ones according to footnote 13. Finally, $n_t^{(1)} = \{0, 1\}$ accounts for the presence or not of time derivatives acting on $G(k_1, \eta')$.

We concentrate now on the time-integral, which can be computed in closed form: by writing $v \equiv n+m+s(+n_t^{(1)}-1)$ and recalling our discussion on the values attainable by n, m, s in section 4.3[16], we find

$$\begin{aligned}\int_{\eta_c}^{0}d\eta'\,\eta'^{-1+v}\,e^{ik_1(1+\cos\theta_j)\eta'} &= \frac{(-1)^v}{(ik_1)^v}\left(\sum_{r=0}^{v-1}\binom{v-1}{r}\frac{r!(-ik_1\eta_c)^{v-1-r}}{(1+\cos\theta_j)^{r+1}}e^{ik_1\eta_c(1+\cos\theta_j)}-\frac{(v-1)!}{(1+\cos\theta_j)^v}\right)\\ &\approx\begin{cases}-\frac{\eta_c^v}{v} & \text{if } |k_1\eta_c(1+\cos\theta_j)|\ll 1\\ \frac{(-1)^{v-1}}{(ik_1)^v}\frac{(v-1)!}{(1+\cos\theta_j)^v} & \text{if } |k_1\eta_c(1+\cos\theta_j)|\gg 1\,.\end{cases}\end{aligned}\quad (65)$$

where in the last passage we have written the leading contribution, also considering the averaging to zero of large oscillations in the final observables.

Inserting this result in equation (64), we obtain the leading contribution

$$\delta_\beta\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\rangle_\lambda^{\rm mv}\approx\mathcal{B}_\lambda^{\rm mv}\,P_{\rm st}(k_1)P_{\rm st}(k_S) \quad (66)$$

with, having substituted back $n+m+s$ in place of v,

$$\mathcal{B}_\lambda^{\rm mv}=\epsilon\,\beta^*_{k_S}\,\frac{\lambda_{nms}}{\sqrt{2\epsilon}}\,\frac{\Lambda M_{\rm Planck}}{H^2}\Big(\frac{k_1}{k_S}\Big)^{1+\min(n,m,s)}\begin{cases}2\big(\frac{H}{\Lambda}|k_S\eta_c|\big)^{n+m+s}\Big(c_{nms}(1)+\frac{c_{nms}(n_t^{(1)})}{(-i|k_1\eta_c|)^{1\text{-}n_t^{(1)}}}\Big) & \text{if } |k_1\eta_c v_{\theta_j}|\ll 1\\ \sum_{j=2}^{3}\Big(\text{-}i\frac{H}{\Lambda}\frac{k_S}{k_1}\Big)^{n+m+s}\Big(\frac{d_{nms}(1)}{v_{\theta_j}^{n+m+s}}+\frac{d_{nms}(n_t^{(1)})}{v_{\theta_j}^{n+m+s+n_t^{(1)}\text{-}1}}\Big) & \text{if } |k_1\eta_c v_{\theta_j}|\gg 1\end{cases} + \text{complex conjugate}\quad (67)$$

where

$$v_{\theta_j}\equiv(1+\cos\theta_j),\qquad c_{nms}(x)\equiv\tfrac{c_\lambda}{2^x(n+m+s-1+x)},\qquad d_{nms}(x)\equiv\text{-}(\text{-}\tfrac{1}{2})^x(n+m+s-2+x)!d_\lambda,\quad (68)$$

and c_λ, d_λ are order one numerical factors, possibly depending on θ_j.

We observe that the generic contributions present both enhancement factors (such as $|k_S\eta_c|\gg 1$, or $\frac{k_S}{k_1}\gg 1$) as well as suppressing ones (such as $\big(\frac{H}{\Lambda}\big)^{n+m+s-1}$ and λ_{nms}). We also observe that for $|k_1\eta_c|\gg 1$ the contribution will not be in general of the local form. However, such contributions could be suppressed and thus subleading. We will discuss enhancements and scale dependence of the results in details in section 5. Observe that for the couplings (60), (61), equation (67) leads to the results we have found before, with the proper c_λ, d_λ, see equations (33), (52).

---

[16] Recall that we are not considering the terms with $n = m = s = 0$ because they are subdominant in slow-roll, that couplings with only two spatial derivatives are reabsorbed by field redefinitions, while those with only one derivative of any type are not present because of isotropy, and also that in the case of time derivatives their order is at most 1 (higher orders must be eliminated using the equations of motions, see [28]).

4.3.2 Modified dispersion relations, Lorentz broken

As discussed in section 4.1.3 the leading contribution to the bispectrum arises when the Whightman functions have support mostly in the interval of times in regions IV/III, given the solution (10). The Whightman functions go as, see appendix A.2,

$$(\partial_{\eta'} G^{\pm})_k(0,\eta') = \text{-}i\frac{H^3}{\dot{\phi}^2} k \frac{(\gamma^{(*)}) e^{\pm i \frac{\Lambda}{H}\Omega(k,\eta')}}{a(\eta') 2k^2}, \quad (\partial_i^{r\geq 0} G^{\pm})_k(0,\eta') = \frac{H^3}{\dot{\phi}^2} (\pm i k_i)^r \frac{\chi^{(*)}(k,\eta')}{a(\eta')} \frac{e^{\pm i\frac{\Lambda}{H}\Omega(k,\eta')}}{2k^2}, \tag{69}$$

where $\chi^{(*)}(k,\eta')$ is defined in equation (103).

Once again, it is convenient to write the contribution to the bispectrum in terms of the variables defined in equation (34). The leading contribution then reads

$$\begin{aligned} \langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle_{\lambda}^{\text{mdr}} &\propto \frac{\lambda_{nms}}{\Lambda^{n+m+s-1}} \sum_{j=2}^{3} \beta^*_{k_j} \frac{H^{n+m+s+5}}{\dot{\phi}^3 2^3 k_1^3 k_2^3 k_3^3} \left(\frac{\Lambda}{H}\right)^{n+m+s} \frac{k_a^{n+1} k_b^{m+1} k_c^{s+1}}{k_S^{n+m+s}} \\ &\times \int_{y_{\rm II}\sim -1}^{y \lesssim \text{-}\frac{H}{\Lambda}} dy' \, (\text{-}y')^{n+m+s-1} \, (i\gamma^{(*)}(x_{a,b,c}, y'))^{n_t^{(a,\,b,\,c)}} (\chi^{(*)}(x_1, y'))^{1\text{-}n_t^{(1)}} e^{i\frac{\Lambda}{H} x_1 \tilde{v}^{(n)}_{\theta_j}(y')} + \text{hermitian}, \end{aligned} \tag{70}$$

where we have used equation (38), $n_t^{(a,\,b,\,c)} = \{0, 1\}$ accounts for the presence or not of time derivatives in the coupling acting on $\zeta_{k_{a,b,c}}$, and a, b, c run over $1, 2, 3$ taking into account the permutations, from which we take the dominant ones according to footnote (13). Now, depending on the specific Lorentz-broken model, n, m, s can assume values larger than 1 ($n + m + s$ larger than 3 [17]).

The integral can be again obtained by asymptotic techniques so that the leading contribution reads

$$\delta_\beta \langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle_{\lambda}^{\text{mdr}} \approx \mathcal{B}_{\lambda}^{\text{mdr}} \, P_{\text{st}}(k_1) P_{\text{st}}(k_S) \tag{71}$$

where, having written $n + m + s = v$ and used table 1 in appendix A.3,

$$\begin{aligned} \mathcal{B}_{\lambda}^{\text{mdr}} &= \epsilon \, \beta^*_{k_S} \frac{\lambda_{nms}}{\sqrt{2\epsilon}} \frac{\Lambda M_{\text{Planck}}}{H^2} \Big(\frac{k_1}{k_S}\Big)^{1+\min(n,m,s)} \\ &\times \sum_{j=2}^{3} \left\{ \begin{array}{ll} \big(\frac{i}{2}\big)^{n_t^{(1)}} \Big(\mathcal{O}(1) + \big(\frac{\Lambda}{H}\frac{k_1}{k_S}\big)^{n_t^{(1)}-1} \mathcal{O}(1)\Big) & \text{if } \frac{\Lambda}{H}\frac{k_1}{k_S} \ll 1 \\ \left[\begin{array}{ll} \big(\frac{H}{\Lambda}\frac{k_S}{k_1}\big)^{v} \Big(\frac{(\text{-}i)^v \, \Gamma(v)}{(1+\cos\theta_j)^v} \big(1 + C^{(0)}_{\theta_j}\big(\frac{H}{\Lambda}\frac{k_S}{k_1}\big)\big) + \mathcal{O}(1)\Big) & \text{if } v_{\theta_j} > \frac{H}{\Lambda}\frac{k_S}{k_1} \\ \big(\frac{H}{\Lambda}\frac{k_S}{k_1}\big)^{\frac{v}{\kappa+1}} \frac{\Gamma(\frac{v}{\kappa+1}) e^{i\frac{\pi v}{2}\frac{\text{sign}(F^{(\kappa)})}{(\kappa+1)}}}{(\kappa+1) |F^{(\kappa)}|^{\frac{v}{\kappa+1}}} \, \big(1 + C^{(\kappa)}_{\theta_j}\big(\frac{H}{\Lambda}\frac{k_S}{k_1}\big)\big) & \text{if } v_{\theta_j} < \frac{H}{\Lambda}\frac{k_S}{k_1} \end{array} \right] & \text{if } \frac{\Lambda}{H}\frac{k_1}{k_S} \gg 1 \end{array} \right\} + \begin{array}{l}\text{complex} \\ \text{conjugate}\end{array} \end{aligned} \tag{72}$$

where

$$C^{(\rho)}_{\theta_j}\Big(\frac{H}{\Lambda}\frac{k_S}{k_1}\Big) \equiv \sum_{r,n=0}^{\infty} c_{r,n} \Big(\frac{H}{\Lambda}\frac{k_S}{k_1}\Big)^{\frac{\kappa+r+n}{\rho+1}} \frac{\Gamma(\frac{v+\kappa+r+n}{\rho+1}) e^{i\frac{\pi(\kappa+r+n)}{(\rho+1)}\frac{\text{sign}(F^{(\rho)})}{2}}}{\Gamma(\frac{v}{\rho+1}) |F^{(\rho+n)}|^{\frac{\kappa+r+n}{\rho+1}}} \tag{73}$$

contains all the subleading asymptotic contributions (the $c_{r,n}$ are calculable order one coefficients, whose precise form is irrelevant for us, obtained from the expansion of the integrand in (70)).

This result is very similar to that obtained in the modified vacuum scenario, except that now n, m, s can also assume values larger than 1 in the case of couplings deriving from the Lorentz-breaking terms in the action, and that $\kappa \to \infty$ for the folded configuration in the modified vacuum scenario, whereas here, for the case of modified dispersion relations, the contribution is again determined at leading order by the first correction to the standard dispersion relations (see (40)). A part from this aspect, the same comment as the end of the previous section applies. The similar structure of the two results can be seen as showing that indeed the squeezing limit is dominated, at leading order, by the generic features common to both scenarios (such as particle creation, interference effects).

---

[17] However, time derivatives are at most of order 1, see footnote 16.

For the couplings (60), (61), equation (72) leads to the results we have found before, see equations (42), (54). Note, in particular, that since the one in equation (43) is the sum of two elementary cubic couplings (one with only time derivatives, the other with one time and two space derivatives) such that for the folded configurations the leading contribution is cancelled out, in those cases the result (54) is given by the first subleading correction in equation (72).

5 Signatures of very high energy physics in the squeezed limit

We will now comment and discuss the results found so far. For observational purposes (regarding the halo bias, for example), the most interesting features of the squeezing limit of the bispectrum are its dependence on the large probed scale k_1^{-1} and the magnitude of the non-Gaussianities [7,8]: therefore, we will focus on those aspects.

From our analysis, it appears that the squeezed limit of the bispectrum shows different kinds of behaviour regarding k_1-dependence and amplitude, subject to the interplay between the squeezed momentum scale k_1 and the scale/time Λ/η_c of new physics. In particular, the different behaviours occur depending on whether $|k_1\eta_c|$ or $\frac{k_1}{k_S}\frac{\Lambda}{H}$ are larger or smaller that 1.

We will momentarily investigate what the different modified scenarios predict concerning the value of $|k_1\eta_c|$ or $\frac{k_1}{k_S}\frac{\Lambda}{H}$, but let us postpone this question for a moment, and discuss first in more details what (and really if) significant enhancements could appear in any case, and what would be the dependence on k_1 for the bispectrum in the squeezing limit.

5.1 Enhancements

We start by discussing the enhancements. Since we have adopted a phenomenological approach we do not have the full-detailed knowledge of the magnitude of β_{k_S}, which the leading contributions to the bispectrum depend upon. However, we know the constraints that the Bogoljubov coefficient has to satisfy for phenomenological reasons and self-consistency of the theory (see section 3.4), and thus we can estimate the largest possible enhancements. Observe also that the leading contribution depends only on β_{k_S}, which does depend on the scale k_S, but not on k_1, thus the k_1-dependence of the leading contribution is not affected by our ignorance of β_{k_S}.

Cases $|k_1\eta_c| \ll 1$ ***and*** $\frac{k_1}{k_S}\frac{\Lambda}{H} \ll 1$

In this case the correction to the bispectrum has a parameter[18] $f_{NL} \sim \mathcal{B}^{\text{mv, mdr}}_{\lambda}$, where

$$|\mathcal{B}^{\text{mv, mdr}}|_\lambda \sim \epsilon\,|\beta_{k_S}|\,\frac{\lambda_{nms}}{\sqrt{2\epsilon}}\,\frac{\Lambda M_{\text{Planck}}}{H^2}\Big(\frac{k_1}{k_S}\Big)^{1+\min(n,m,s)}\Big(\frac{H\,D}{\Lambda}\Big)^{n+m+s}\Big(1+\Big(\frac{k_1 D}{k_S}\Big)^{n_t^{(1)}\text{-}1}\mathcal{O}(1)\Big) \tag{74}$$

$$\leq \epsilon\sqrt{|\mu|}\,\frac{\lambda_{nms}}{\sqrt{2}}\,\frac{M^2_{\text{Planck}}}{H\Lambda}\Big(\frac{k_1}{k_S}\Big)^{1+\min(n,m,s)}\Big(\frac{H\,D}{\Lambda}\Big)^{n+m+s}\Big(1+\Big(\frac{k_1 D}{k_S}\Big)^{n_t^{(1)}\text{-}1}\mathcal{O}(1)\Big)\,. \tag{75}$$

In the inequality (75) we have used the constraint (16) on $|\beta_{k_S}|$. Here, $D = |k_S\eta_c|$ in the modified vacuum case, while $D = \frac{\Lambda}{H}$ in the scenario with modified dispersion relations[19]. The similarity of the results in the two different scenarios is an indication that the squeezing limit is dominated by the generic features common to both of them (particle creation, interference effects).

---

[18] As before, 'mv' indicates the modified vacuum scenario, while 'mdr' the one with modified dispersion relations.

[19] However, in the NPHS approach $|k_S\eta_c| = \frac{\Lambda}{H}$.

If we consider first $D=|k_S\eta_c|$, the inequality reads

$$|\mathcal{B}^{D=|k_S\eta_c|}|_\lambda \leq \epsilon\sqrt{|\mu|}\,\frac{\lambda_{nms}}{\sqrt{2}}\,\frac{M^2_{\text{Planck}}}{H\Lambda}\Big(\frac{k_1}{k_S}\Big)^{1+\min(n,m,s)}\Big(\frac{|p_S(\eta_c)|}{\Lambda}\Big)^{n+m+s}\Big(1+\Big(\frac{|p_1(\eta_c)|}{k_S}\Big)^{n_t^{(1)}\text{-}1}\mathcal{O}(1)\Big)\,, \tag{76}$$

where $p_{S(1)}(\eta_c)=-k_{S(1)}\eta_c H$ is the physical momentum associated with the wavenumber $k_{S(1)}$ at the boundary time η_c.

When instead $D=\frac{\Lambda}{H}$, such as for the cases of NPHS vacuum or modified dispersion relations, the expression for $|\mathcal{B}|_\lambda^{\text{NPHS, mdr}}$ simplifies and we obtain

$$|\mathcal{B}^{\text{NPHS, mdr}}|_\lambda \leq \epsilon\sqrt{|\mu|}\,\frac{\lambda_{nms}}{\sqrt{2}}\,\frac{M^2_{\text{Planck}}}{H\Lambda}\Big(\frac{k_1}{k_S}\Big)^{1+\min(n,m,s)}\Big(1+\Big(\frac{k_1}{k_S}\frac{\Lambda}{H}\Big)^{n_t^{(1)}\text{-}1}\mathcal{O}(1)\Big)\,. \tag{77}$$

From (76), (77), we can understand various characteristics of the amplitude of the non-Gaussianities when $|k_1\eta_c|$, $\frac{k_1}{k_S}\frac{\Lambda}{H}\ll 1$. First of all, we see from (76) that the maximum amplitude for modified vacuum scenarios occurs when the physical momentum $p_S(\eta_c)$ associated with k_S at the particle creation time η_c is really at the largest scale that we can trust our effective theory at: the scale of new physics Λ. But this is precisely what occurs in the NPHS scenario of vacuum modification, leading to (77). In the case of modified dispersion relations, equation (77) arises for the same reason: the physical momentum that would appear in place of $p_S(\eta_c)$ in the analogous of (76) is $p_S(\eta_{\text{II}}^{(k_S)})$, at the particle creation/WKB breaking time $\eta_{\text{II}}^{(k_S)}$, which has in fact a magnitude of the order of Λ.

However, in order to see if there can be actual overall enhancements, potentially interesting for observations, one needs to take into account also the other factors in the equations for $\mathcal{B}_\lambda^{\text{mv, mdr}}$: both the suppressing ($\lambda_{nms}\,\epsilon\sqrt{|\mu|}\,(\frac{k_1}{k_S})^{1+\min(n,m,s)}$), and the enhancing ($\frac{M^2_{\text{Planck}}}{H\Lambda}$). In this respect, it is immediately evident that terms with higher number of derivatives are more suppressed (since $\min(n,m,s)$ will be higher). This is particularly true for the Lorentz-breaking couplings in the scenarios with modified dispersion relations, where n,m,s can assume values even much higher that one.

As it was in fact imaginable, the cubic couplings (22) and (43) that we have discussed as detailed examples, which have the coupling constants and order of derivatives as summarized in equations (60), (61), provide (among) the least suppressed contributions, with $\big(\frac{k_1}{k_S}\big)^0$ and $\big(\frac{k_1}{k_S}\big)^2$ suppression respectively (a part from the suppressing factors $\lambda_{11\text{-}1},\lambda_{111}$, $\epsilon\sqrt{|\mu|}$, of course). The minimal cubic coupling is in fact also leading in the standard scenario, see [6].

If we concentrate on these dominant (less suppressed) couplings, which we have studied in details, we find that there can be actual overall enhancement of the non-Gaussianities, with respect to the standard scenario, for certain values of the scale of new physics. Let us discuss a concrete example: take $\epsilon\sim|\mu|\sim 10^{-2}$, $H\sim 10^{-5}M_{\text{Planck}}$, compatible with the results of WMAP, and $\Lambda\sim 10^{-3}M_{\text{Planck}}$, which corresponds to the supersymmetric GUT scale. Then, if we can probe down to $\frac{k_1}{k_S}\lesssim 10^{-2}$, the largest possible enhanced amplitude with respect to the standard scenario that we could observe would be roughly of the order of

$$|\mathcal{B}^{\text{NPHS, mdr}}|_{\text{min cub}}\sim 10\,\epsilon\sim 10\,(1-n_s)\,. \tag{78}$$

in the case of the minimal coupling (60), while

$$|\mathcal{B}^{\text{NPHS, mdr}}|_{\text{quart der}}\sim 10^3 g\,(1-n_s)\sim 10\,(1-n_s) \tag{79}$$

in the case of the quartic derivative interaction (61) with $g\sim 10^{-2}$. It is notable how, for the higher (quartic)-derivative interaction, the additional suppressing factor $\big(\frac{k_1}{k_S}\big)^2$, and enhancing factor $\frac{M^2_{\text{Planck}}}{\Lambda^2}$ (essentially due to the power of a^{-1} scaling) compensate each other to give a result of the same magnitude as that of the minimal cubic coupling: a result which is quite different from the one in the standard scenario (where the a^{-1} scaling of the interactions does not have any important effect, since the dominant contributions to non-Gaussianities occur at late times).

Observe that, according to the analysis of [6], values such as $\frac{k_1}{k_S} \sim 10^{-2}$ are indeed within the reach of LSS investigations, such as for example EUCLID, which in the most optimistic estimates can probe values down to $\frac{k_1}{k_S} < 10^{-3}$ [6].

Cases $|\mathbf{k_1}\eta_c| \gg \mathbf{1}$ ***and*** $\frac{\mathbf{k_1}}{\mathbf{k_S}}\frac{\mathbf{\Lambda}}{\mathbf{H}} \gg 1$

In this case the correction to the bispectrum has a parameter $f_{NL} \sim \mathcal{B}_g^{\text{mv, mdr}}$ with

$$|\mathcal{B}^{\text{mv, mdr}}|_\lambda \sim \epsilon\,|\beta_{k_S}|\,\frac{\lambda_{nms}}{\sqrt{2\epsilon}}\,\frac{\Lambda M_{\text{Planck}}}{H^2}\,(\frac{k_1}{k_S})^{1+\min(n,m,s)}E^{(n,m,s)}(\frac{H}{\Lambda}\frac{k_S}{k_1}) \tag{80}$$

$$\leq \epsilon\sqrt{|\mu|}\;\frac{\lambda_{nms}}{\sqrt{2}}\;\frac{M^2_{\text{Planck}}}{H\Lambda}\,(\frac{k_1}{k_S})^{1+\min(n,m,s)}E^{(n,m,s)}(\frac{H}{\Lambda}\frac{k_S}{k_1}) \tag{81}$$

where in the last inequality we have used the constraint (16) on $|\beta_{k_S}|$. Here,

$$E^{(n,m,s)}(\frac{H}{\Lambda}\frac{k_S}{k_1}) \simeq \begin{cases} \left(\frac{H}{\Lambda}\frac{k_S}{k_1}\right)^{n+m+s}\Big(\mathcal{O}(1)+\sum_{r=0}^{\infty}\mathcal{O}\big((\frac{H}{\Lambda}\frac{k_S}{k_1})^{\kappa+r}\big)\Big) & \text{if } 1+\cos(\theta_j) > \frac{H}{\Lambda}\frac{k_S}{k_1} \\ \begin{bmatrix} \big(\frac{H\,D}{\Lambda}\big)^{n+m+s}\Big(1+\big(\frac{k_1 D}{k_S}\big)^{n_t^{(1)}\text{-}1}\mathcal{O}(1)\Big) & \text{mv} \\ \big(\frac{H}{\Lambda}\frac{k_S}{k_1}\big)^{\frac{n+m+s}{\kappa+1}}\Big(1+\sum_{r=0}^{\infty}\mathcal{O}\big((\frac{H}{\Lambda}\frac{k_S}{k_1})^{\frac{\kappa+r}{\kappa+1}}\big)\Big) & \text{mdr} \end{bmatrix} & \text{if } 1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{k_S}{k_1} \end{cases} \tag{82}$$

Since the condition leading to this result is $\frac{k_1}{k_S}\frac{\Lambda}{H} \gg 1$, the powers of $\big(\frac{H}{\Lambda}\frac{k_S}{k_1}\big)$ in equations (81), (82) will be smaller than one, and thus not an enhancing factor. From (82) it appears that in the scenario of modified dispersion relations, the (nearly) folded configurations, singled out by the condition $1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{k_S}{k_1} \ll 1$, are enhanced with respect to the other configurations, but not as much as in the modified vacuum scenario. In any case. the result shows a non-trivial θ_j dependence (recall that we consider a realistic squeezing limit where k_1 is small but non-zero). Furthermore, we see that the higher the number of derivatives, the more the suppression because of the factor $\big(\frac{k_1}{k_S}\big)^{1+\min(n,m,s)}$: again, the cubic couplings (22) and (43) that we have discussed as detailed examples turn out to be leading.

Equation (81) also includes enhancement factors such as $\frac{M^2_{\text{Planck}}}{H\Lambda}$, thus, once more, the possibility or not of actual overall enhancements depends on the values of the scales and quantities at play. Let us again consider the same example as before and the couplings (22) and (43), to give some pointers about the possibilities. Recall that we take $\epsilon \sim |\mu| \sim 10^{-2}$, $H \sim 10^{-5}M_{\text{Planck}}$, compatible with the results of WMAP, and $\Lambda \sim 10^{-3}M_{\text{Planck}}$ (supersymmetric GUT scale). Then, if we can only probe values down to $\frac{k_1}{k_S} \gtrsim 10^{-2}$, we are in the case $\frac{k_1}{k_S}\frac{\Lambda}{H} > 1$ and we find the bound, roughly,

$$|\mathcal{B}^{\text{mv, mdr}}|_{\text{min cub}} \lesssim \begin{cases} 10^{-1}\frac{k_S}{k_1}(1-n_s) & \text{if } 1+\cos(\theta_j) > \frac{H}{\Lambda}\frac{1}{x_1} \\ \begin{bmatrix} 10(1-n_s) & \text{mv} \\ 10^{\frac{\kappa\text{-}1}{\kappa+1}}\big(\frac{k_S}{k_1}\big)^{\frac{1}{\kappa+1}} & \text{mdr} \end{bmatrix} & \text{if } 1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{1}{x_1} \end{cases}\;. \tag{83}$$

We see that for $\frac{k_1}{k_S} \simeq 10^{-2}$ the folded configurations, both in the modified vacuum and modified dispersion relations scenarios, could be ten time enhanced with respect to the standard result, whereas the other configurations would present only a moderate enhancement compared to the result in the standard scenario.

In the case of the higher-derivative interaction (43), for a coupling $g \sim 10^{-2}$, we find, see (52), (54) and the comment at the end of section 4.3.2, after (72),

$$|\mathcal{B}^{\text{mv, mdr}}|_{\text{quart der}} \lesssim \begin{cases} 10^{-1}\frac{k_S}{k_1}(1-n_s) & \text{if } 1+\cos(\theta_j) > \frac{H}{\Lambda}\frac{1}{x_1} \\ \begin{bmatrix} 10^3\left(\dfrac{k_1}{k_S}\right)(1-n_s) & \text{mv} \\ 10^{3-\frac{4}{\kappa+1}}\left(\dfrac{k_S}{k_1}\right)^{\frac{2}{\kappa+1}-1}(1-n_s) & \text{mdr} \end{bmatrix} & \text{if } 1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{1}{x_1} \end{cases} . \tag{84}$$

For $\frac{k_1}{k_S} \simeq 10^{-2}$, (83) and (84) are of the same magnitude, which again confirms that in the modified scenarios the different a^{-1} scaling of the higher-derivative interactions can compensate for the stronger suppression by higher powers of $\frac{k_1}{k_S}$ and make the squeezing limit sensitive to higher-derivative interactions.

5.2 Dependence on k_1

We discuss now the dependence of the corrections to the bispectrum in the squeezing limit on the squeezed wavenumber k_1. Recall that although β_{k_S} depends on the scale k_S in a way that we cannot fully determine with our phenomenological approach, it does not depend on k_1, thus the k_1-dependence of the leading correction to the bispectrum is not affected by our ignorance of the scale dependence of β_{k_S}.

In the following, we will rewrite equations (66)-(67) and (71)-(72) highlighting the dependence on k_1. Again, we find that the field theoretical results for the modified scenarios have two different kinds of behaviour, depending on the magnitude of the scale of new physics as compared to the sensitivity of the observations. We will show that the results are different from the one in the standard scenario, which grows as $\sim k_1^{-3}$ in the squeezing limit [4–6], and from the one obtained with the folded template proposed in [9] to model modified vacuum scenarios, which yields a bispectrum growing as $\sim k_1^{-2}$ [7,8].

Cases $|k_1\eta_c| \ll 1$ ***and*** $\frac{k_1}{k_S}\frac{\Lambda}{H} \ll 1$

In this case, equations (66)-(67) and (71)-(72) yield

$$\delta_\beta\langle\zeta_{\vec{k}_1}(\eta)\zeta_{\vec{k}_2}(\eta)\zeta_{\vec{k}_3}(\eta)\rangle_\lambda \approx \epsilon\,|\beta_{k_S}|\,\frac{\lambda_{nms}}{\sqrt{2\epsilon}}\,\frac{M^2_{\text{Planck}}}{H\Lambda}\left(\frac{HD}{\Lambda}\right)^{n+m+s}\left(\frac{k_1}{k_S}\right)^{1+\min(n,m,s)}\left(1+c_{n_t^{(1)}}\left(\frac{k_1 D}{k_S}\right)^{n_t^{(1)}-1}\right)P_{\text{st}}(k_1)P_{\text{st}}(k_S), \tag{85}$$

with $D = |k_S\eta_c|$ or $\frac{\Lambda}{H}$, respectively for modified vacuum or modified dispersion relations.

The correction scales as $k_1^{-3+\min(n,m,s)+n_t^{(1)}}$, $n_t^{(1)} = \{0,1\}$. We thus find that higher-derivative couplings (larger values for n,m,s) grow more slowly. The fastest growth for small k_1 is of the form

$$\delta_\beta\langle\zeta_{\vec{k}_1}(\eta)\zeta_{\vec{k}_2}(\eta)\zeta_{\vec{k}_3}(\eta)\rangle_\lambda \approx \epsilon\,|\beta_{k_S}|\,\frac{\lambda_{nms}}{\sqrt{2\epsilon}}\,\frac{M^2_{\text{Planck}}}{H\Lambda}\left(\frac{H\,D}{\Lambda}\right)^{n+m+s} P_{\text{st}}(k_1)P_{\text{st}}(k_S)\,, \tag{86}$$

that is, a local shape k_1^{-3}, and can be found for couplings such as the minimal coupling cubic, with $\min(n,m,s) = -1$ (recall that this value occurs due to the combination $(\vec{\partial})^{-2}\dot{\varphi}$, therefore it is always accompanied by $n_t^{(1)} = 1$).

Also in these cases, however, although it has a local form, the result is different from the one in the standard scenario because non-Gaussianities can be enhanced.

Cases $|k_1\eta_c| \gg 1$ ***and*** $\frac{k_1}{k_S}\frac{\Lambda}{H} \gg 1$

In this case, equations (66)-(67) and (71)-(72) yield, at leading order,

$$\delta_\beta \langle \zeta_{\vec{k}_1}(\eta)\zeta_{\vec{k}_2}(\eta)\zeta_{\vec{k}_3}(\eta)\rangle_\lambda \approx \epsilon\,|\beta_{k_S}|\,\frac{\lambda_{nms}}{\sqrt{2\epsilon}}\,\frac{M^2_{\rm Planck}}{H\Lambda}$$
$$\times \Big(\frac{k_S}{k_1}\Big)^{-1-\min(n,m,s)} \left\{\begin{matrix} \Big(\frac{H}{\Lambda}\frac{k_S}{k_1}\Big)^{n+m+s} & \text{if } 1+\cos(\theta_j) > \frac{H}{\Lambda}\frac{1}{x_1} \\ \begin{bmatrix} \Big(\frac{H\,D}{\Lambda}\Big)^{n+m+s}\Big(1+c_{n_t^{(1)}}\Big(\frac{k_1 D}{k_S}\Big)^{n_t^{(1)}\text{-}1}\Big) & \text{mv} \\ \Big(\frac{H}{\Lambda}\frac{k_S}{k_1}\Big)^{\frac{n+m+s}{\kappa+1}} & \text{mdr} \end{bmatrix} & \text{if } 1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{1}{x_1} \end{matrix}\right\} P_{\rm st}(k_1)P_{\rm st}(k_S)\,. \tag{87}$$

Except for the case of folded configurations in modified vacuum scenarios, to which the same considerations regarding equation (85) apply, it appears that higher-derivative couplings (larger values for n, m, s) would lead to a more pronounced growth for small k_1 as its exponent will be more negative. However, all these contributions are very suppressed because of the factors $\big(\frac{H}{\Lambda}\big)^{n+m+s}$, $\big(\frac{H}{\Lambda}\big)^{\frac{n+m+s}{\kappa+1}}$. The most interesting cases are thus those that are least suppressed, like the minimal cubic coupling that we studied as a detailed example, where $n+m+s=1$ and $\min(n,m,s)=-1$, see (60), in which case

$$\delta_\beta \langle \zeta_{\vec{k}_1}(\eta)\zeta_{\vec{k}_2}(\eta)\zeta_{\vec{k}_3}(\eta)\rangle_{\rm min\ cub} \approx \epsilon\,|\beta_{k_S}|\,\frac{M_{\rm Planck}}{\Lambda} \left\{\begin{matrix} \frac{H}{\Lambda}\frac{k_S}{k_1} & \text{if } 1+\cos(\theta_j) > \frac{H}{\Lambda}\frac{1}{x_1} \\ \begin{bmatrix} 1 & \text{mv} \\ \Big(\frac{H}{\Lambda}\frac{k_S}{k_1}\Big)^{\frac{1}{\kappa+1}} & \text{mdr} \end{bmatrix} & \text{if } 1+\cos(\theta_j) < \frac{H}{\Lambda}\frac{1}{x_1} \end{matrix}\right\} P_{\rm st}(k_1)P_{\rm st}(k_S)\,. \tag{88}$$

We see that in this case the form of the bispectrum in the squeezing limit is *i*) not local, with a scaling k_1^{-4}, and with amplitude not too severely suppressed for non-folded configurations, *ii*) enhanced (see also section 5.1), with a non-local shape for (nearly) folded configurations in the case of modified dispersion relations, and *iii*) with a local shape and greater enhancement for folded configurations in the modified vacuum case. Remarkably, the scaling with k_1 of the contribution from (nearly) folded configurations in the case of modified dispersion relation captures the power κ of the first momentum correction to the standard dispersion relation.

5.3 Predictions from the different scenarios

We have seen that, most interestingly, in modified scenarios the leading features of the bispectrum in the squeezing limit depend on very general aspects of the theory (exit from horizon, particle production, interference). These cannot determine the bispectrum in the squeezing limit in all details, but even adopting a wide-range phenomenological approach, we have been able to determine both the leading dependence on the squeezed scale k_1 and the presence of and bounds on enhancements of non-Gaussianities.

We have seen that these features depend on the values (less or more than 1) of the quantities $|k_1\eta_c|$, $\frac{k_1}{k_S}\frac{\Lambda}{H}$. We would like now to ask ourselves which kind of conditions or predictions the different theoretical scenarios impose or make about the values of $|k_1\eta_c|$ and $\frac{k_1}{k_S}\frac{\Lambda}{H}$, and, thus, what features of the bispectrum would be realized in the squeezing limit.

- *EBA modified vacuum.* In the Effective Boundary Action approach, the boundary condition for the fields is fixed at a time η_c independently of the modes k ("beginning of inflation"). In that scenario, it is then assumed that all modes probed by observations were subhorizon at η_c, because in that case one can fix initial conditions with more confidence. Therefore, since also k_1 is an observable mode, it must be $|k_1\eta_c| > 1$ (subhorizon mode at η_c).

Therefore, if the EBA is the correct way to model the effects of very high energy physics from a low-energy point of view encoding them in an initial state for the perturbations, then the leading correction to the bispectrum for non-folded configurations would grow at large scales as k_1^{-4}, but with only moderately enhanced non-Gaussianities, whereas the contribution from folded configurations would be more enhanced, but growing only as k_1^{-3}, see (80), (83), (87), (88).

- *NPHS modified vacuum.* In the NPHS approach, instead, the boundary conditions is imposed at the time given by $|k\,\eta_c(k)| = \frac{\Lambda}{H} \gg 1$. In this case, the picture is that different modes are created out of the vacuum at different times in a certain condition that fixes the vacuum of the effective theory at scales lower than Λ. When computing the three-point function, therefore, we cannot trace the interaction of the modes backward in time for a time earlier that the "creation time" for the largest of the considered modes: the prior evolution (and our ignorance about it) is encoded in the initial state. Thus, in the case of the squeezed limit, the earliest time possible is related to the large momentum scale $k_{\rm max} \equiv \max(k_{1,2,3}) \simeq k_S$, and not to k_1, so that $\frac{k_1}{k_S}\frac{\Lambda}{H}$ can be both larger or smaller than 1, depending on the values of the scales at play in the specific model.

  In particular, the leading correction to the bispectrum would grow at large scales as k_1^{-4} for non-folded configurations, only provided that the scale of new physics is such that $\Lambda > \frac{k_S}{k_1}H$, where $\frac{k_1}{k_S}$ is the smallest ratio of scales that can be probed, see (87), (88). For folded configurations, and when $\Lambda < \frac{k_S}{k_1}H$, the contribution would instead be of the local type, growing at most as k_1^{-3}.

  An interesting outcome of this result, is that in case we could detect one or the other behaviour, say in the halo bias, and separate it from other sources of non-Gaussianities, we could obtain information on the magnitude of Λ, given our knowledge of the observation sensitivity $\frac{k_1}{k_S}$.

- *Modified dispersion relations.* In the scenarios with modified dispersion relations violating WKB at early times, in case of a Lorentz-breaking scale Λ such that $\frac{k_1}{k_S}\frac{\Lambda}{H} > 1$, where $\frac{k_1}{k_S}$ is the smallest ratio that can be probed, there would be contributions to the bispectrum growing as fast as k_1^{-3-N} with N only bound by the specific model, and in principle also possibly large, due to the presence of higher-derivative Lorentz-breaking couplings, see (87). However, we have found that those contributions are severely suppressed by powers $\left(\frac{H}{\Lambda}\right)^M$, $M \geq N$. The least suppressed contributions (which could actually be enhanced) show at leading order a growth as k_1^{-4} for non-unfolded configurations, and slower powers k_1^{-r}, with r fractional, for the (nearly) folded ones, see (87), (88). Remarkably, in the latter case, the exponent of k_1 captures, being directly related to it, the momentum power of the lowest correction to the standard dispersion relation.

  On the other hand, for $\Lambda < \frac{k_S}{k_1}H$, the leading contribution would be of the local type growing at most as k_1^{-3}, see (86). Once again, if we could observe one or the other behaviour, we would obtain information on the magnitude of the Lorentz-breaking scale from our knowledge of the probed k_1.

6 Comments on the halo bias

We will here briefly comment on the implications of our results for the halo bias, leaving a more detailed analysis for future research. The halo bias depends on the bispectrum and the spectrum as [7] ([8] has a similar but not identical formula)

$$\frac{\Delta b_h(k_1,R)}{b_h} = \frac{\delta_c}{D(z)\mathcal{M}_R(k_1)}\frac{1}{8\pi^2\sigma_R^2}\int_0^\infty dk_2\,k_2^2\mathcal{M}_R(k_2) \times\int_{-1}^{1} d\xi\,\mathcal{M}_R\big(\sqrt{k_1^2+k_2^2+2k_1k_2\xi}\big)\frac{B(k_2,\sqrt{k_1^2+k_2^2+2k_1k_2\xi},k_1)}{P(k_1)}\,, \quad (89)$$

where σ_R^2 is the variance of the dark matter density perturbations smoothed on a scale of Lagrangian radius R, δ_c is the critical threshold for the collapse of a spherical object (for a matter-dominated universe $\delta_c = 1.686$), $D(z)$ is the linear growth factor normalized to be $D(z) = (1+z)^{-1}$ during matter domination, P and B are the two-point correlator[20] and bispectrum for ζ, and $\mathcal{M}_R$ is the linear relation between the dark matter density perturbations smoothed on a scale R and the primordial curvature perturbation:

$$\mathcal{M}_R(k) \equiv \frac{2k^2}{5\Omega_m H_0^2} T(k) W_R(k) \,, \tag{90}$$

where Ω_m is the present time fractional density of matter. Finally, H_0 is the present Hubble rate, T is the transfer function normalized to one on large scales and W_R is the filter function with characteristic scale R.

The halo bias depends on both the primordial bispectrum and the subsequent gravitational processing which generates additional non-Gaussianities. These General Relativity corrections have been studied for example in [7] and it has been shown that they give a typical contribution to the halo bias, which could permit to distinguish them. We will therefore comment only the primordial contributions.

The behaviour of the halo bias at large scales has been computed for the local, equilateral and folded *templates* for the primordial non-Gaussianities in [7] (see also [8]), finding that the bias goes approximately as k_1^{-2} with the local template, as a constant in k_1 with the equilateral, and as k_1^{-1} with the folded (the folded template goes indeed as k_1^{-2}).

The latter result is the one that can be compared with ours in the case of the modified vacuum approach, since the template was proposed as an approximation to model the bispectrum for modified vacuum scenarios [9]. In fact, we have shown that this leads to the wrong result in the case of the squeezing limit: the template does not depend on the additional scale(s) η_c, Λ and thus its squeezing limit is different than the one that actually occurs in the rigorous field theory computation. We have indeed shown in sections 4.1.2, 4.2.1 and 4.3.2, that for realistic limits (in view of realistic observations), the leading contribution to the bispectrum grows as k_1^{-4} or k_1^{-3}, depending on the compared magnitudes of the scale of new physics and the sensitivity of the observation (smallest squeezed momentum scale), so that the halo bias would approximately grow either as k_1^{-3} or k_1^{-2}, and not as k_1^{-1}.

Similarly, in the scenario of modified dispersion relations violating WKB at early times, the behaviour of the bispectrum in the squeezed limit, see sections 4.1.3, 4.2.2, 4.3.2, would also lead to a bias that approximately grows either as k_1^{-3} or k_1^{-2} for small k_1 for non-folded configurations, again depending on the relative magnitudes of the high-energy scales and the sensitivity of the observations. (Nearly) folded configurations would instead show different scalings with k_1, determined by the exponent of the first correction to the standard dispersion relation for small momenta.

In general, a feature relevant for the observations of the bias is that the high energy modifications could enhance the single-field non-Gaussianities in the squeezing limit, compared to the practically undetectable level that is present in the standard scenarios of single field models with Bunch-Davies vacuum and standard kinetic terms. However, the final magnitude of the non-Gaussianities depends as well on the specific model, which determines the precise form and magnitude of the coefficients β_{k_S} entering the equations.

7 Final discussion and conclusion

In this article we have analized the squeezing limit of the bispectrum in single-fields models of inflation when modifications of the theory at very high energy are present. Due to the large number of inflationary models available, it is useful to look at general features that can be obtained when these modifications are rather generic and simple. In particular, we have considered 1) the presence of terms with higher derivatives, which are not suppressed any more beyond a certain momentum scale and modify the dispersion relations of the fields, and 2) the possibility of initial condition for the solution of the field equation that parametrize in a simple and generic way our ignorance on the physics at very high energies. The leading contribution to the squeezing limit appears dominated by very general features of the scenarios (particle creation at early times, interference) and thus it is possible to obtain general results and bounds.

---

[20] In the literature on the halo bias, the two-point correlator is often called the spectrum [7,8], not to be confused with the quantity $\mathcal{P} = \frac{k^3}{2\pi^2} P$, also called spectrum, usual in the CMBR literature.

Those that we have found are interesting in two respect: on the one hand, they show that non-Gaussianities could be enhanced by high energy modifications of the theory, which would improve the possibility of detection of those phenomena. On the other, the behaviour of the bispectrum at large scale in these scenarios is distinctive and differs from the one that has been found in the standard scenarios, with Bunch-Davies vacuum and standard dispersion relation, and also different from that found in the previous literature using the approximate template proposed in [9] for the modified vacuum approach. Particle creation at early times and interference effects explain the differences with the standard result. Instead, the differences with the result obtained using the template for the modified vacuum approach arise principally because the approximate template does not actually depend on the scale of the new physics, while the bispectrum calculated from the field theory does, and therefore taking the squeezing limit in the template cannot account for the interplay between the scale of the small mode k_1 and that of the new physics when taking the limit itself.

Our results show that it could be possible to obtain various different pieces of information about the features of the high energy physics theory from large scale observations such as those of the halo bias, if the signatures we have found were detected. In fact, the dependence of the bispectrum on the small wavenumber k_1 in the realistically probed squeezing limit changes in relation with the magnitude of the new physics scale: if able to detect certain dependences of the bispectrum, one can place bounds on the new scale of physics.

Of course, other sources could give origin to sizable non-Gaussianities of non-local shape, for example in multi-field models, and thus an interesting question for future research would then be if the effects due to high-energy modifications of the theory that we have considered can be distinguished from those. For example, the dependence on the small mode k_1 of the halo bias could be different, or the enhancement factors could be absent or of different magnitude. We leave this as an interesting outlook for future research.

Acknowledgments

The author is supported by a Postdoctoral F.R.S.-F.N.R.S. research fellowship via the Ulysses Incentive Grant for the Mobility in Science.

A Appendices

A.1 Solution of the field equation in the case of modified dispersion relations

We parametrize the modifications to the dispersion relation via a *generic* function $F\left(\frac{H}{\Lambda}k\eta\right)$:

$$\omega(\eta,k) = a(\eta)\omega_{\text{phys}}(p) = a(\eta)p\,F\left(-\frac{p}{\Lambda}\right) = k\,F\left(\frac{H}{\Lambda}k\eta\right)\,, \qquad F(x\to 0)\to 1\,, \qquad \frac{H}{\Lambda}\ll 1\,, \tag{91}$$

with the only request that the WKB conditions be violated for a short period of time at early times. The generic shape of such a dispersion relation is in figure 1. The approximate solution of the equation of motion (5) in these cases is given in equation (10), where, see [16]:

- in regions I and III, in terms of the convenient variable $\frac{H}{\Lambda}k\eta \equiv y_k$:

$$u_1(y_k) = \frac{e^{-i\frac{\Lambda}{H}\Omega(y_k)}}{\sqrt{2\,k\,U(y_k)}}\,, \qquad u_2(y_k) = u_1^*(y_k) = \frac{e^{+i\frac{\Lambda}{H}\Omega(y_k)}}{\sqrt{2\,k\,U(y_k)}}\,, \tag{92}$$

  with

$$\Omega(y_k) = \int^{y_k} U(y_k')dy_k'\,, \qquad U(y_k) \equiv F + \epsilon^2\left(-\frac{\partial_y^2 F}{4F^2} + 3\frac{(\partial_y F)^2}{8F^3} - \frac{1}{F y_k^2}\right), \tag{93}$$

- in region IV, where $\omega(\eta,k) \sim k$ and $k\eta \lesssim 1$,

$$\mathcal{V}_{1,2} = \sqrt{-\eta}H_\nu^{(1,2)}(-k\eta)\big|_{k\eta\lesssim 1}, \qquad \nu \sim \frac{3}{2} \tag{94}$$

- in region II, the partial solution $\mathcal{U}_{1,2}(\eta,k)$ can be found either by solving in details the specific equation, if one has a favourite model, or, more generally, by expanding the frequency around the minimum ω_0 at $\eta = \eta_m$ as

$$\omega^2 = a^2\omega^2_{\text{phys}} \simeq \frac{1}{\eta^2 H^2}\left(\omega_0^2 + \omega_0\omega''_{\text{phys}}(\eta_m)(\eta-\eta_m)^2\right), \tag{95}$$

  so that

$$\begin{aligned} f_k(\eta)\Big|_{\text{II}} &= \sum_{i=1}^{2} B_i\,\mathcal{U}_i(\eta,k) = B_1\,W(i\kappa\,\eta_m,\sigma,2i\kappa\,\eta) + B_2\,W(-i\kappa\,\eta_m,\sigma,-2i\kappa\,\eta) \\ \kappa &\equiv \frac{\sqrt{\omega_0\omega''_{\text{phys}}(\eta_m)}}{H} \qquad \sigma \equiv \frac{\sqrt{9H^2 - 4\omega_0^2 - 4H^2\kappa^2\eta_m^2}}{2H}\,, \end{aligned} \tag{96}$$

  where $W(a,b,z)$ is the Whittacker function. As explained in [16], this approximation is well-justified also because backreaction constraints the interval $[\eta_{\text{I}},\eta_{\text{II}}]$ to be very small ($\Delta < 1$), so that $\eta \sim \eta_m$ for $\eta,\eta_m \in [\eta_{\text{I}},\eta_{\text{II}}]$. In any case, it actually turns out that the details of the solution in region II are not important for what concerns the leading contribution to the bispectrum [16].

By asking for the continuity of the function and its first derivative, and imposing the Wronskian condition $\mathcal{W}\{f,f^*\} = -i$ to have the standard commutation relations in the quantum theory, we obtain in full generality

$$D_1 = \frac{\sqrt{\pi}}{2}e^{i\frac{\pi}{2}\nu+i\frac{\pi}{4}}\alpha_k \qquad\qquad D_2 = \frac{\sqrt{\pi}}{2}e^{-i\frac{\pi}{2}\nu-i\frac{\pi}{4}}\beta_k \tag{97}$$

$$B_1 = \frac{\mathcal{W}\{\varsigma_k\,u_1,\mathcal{U}_2\}}{\mathcal{W}\{\mathcal{U}_1,\mathcal{U}_2\}}\Bigg|_{\eta_{\text{I}}} \qquad\qquad B_2 = -\frac{\mathcal{W}\{\varsigma_k\,u_1,\mathcal{U}_1\}}{\mathcal{W}\{\mathcal{U}_1,\mathcal{U}_2\}}\Bigg|_{\eta_{\text{I}}} \tag{98}$$

$$\alpha_k = \frac{\mathcal{W}\{B_1\mathcal{U}_1 + B_2\mathcal{U}_2,u_2\}}{\mathcal{W}\{u_1,u_2\}}\Bigg|_{\eta_{\text{II}}} \qquad\qquad \beta_k = -\frac{\mathcal{W}\{B_1\mathcal{U}_1 + B_2\mathcal{U}_2,u_1\}}{\mathcal{W}\{u_1,u_2\}}\Bigg|_{\eta_{\text{II}}}\,, \tag{99}$$

where $\mathcal{W}$ is the Wronskian. We choose $\varsigma_k = 1$ picking up the usual adiabatic vacuum. The Wronskian condition also imposes $\alpha_k^2 - \beta_k^2 = 1$. By expanding for small Δ around η_{I}, we obtain equation (12) in the text.

Finally, the parameter signalling the WKB violation is [16]

$$\mathcal{Q} = -\left(-\frac{\omega''}{4\omega^2} + 3\frac{\omega'^2}{8\omega^3} - \frac{1}{\omega\eta^2}\right)^2 + \frac{\omega''}{2\omega} - 3\frac{\omega'^2}{4\omega^2} - \frac{U''}{2U} + 3\frac{U'^2}{4U^2}\,. \tag{100}$$

A.2 Formulas for Whightman functions and their time derivatives

The bispectrum is computed at a time η when all momenta have exit the horizon, therefore the Whightman functions have the general form

$$G_k(\eta\sim 0,\eta') = \lim_{\eta\to 0}\frac{H^2}{\dot\phi^2}\frac{f_k(\eta)}{a(\eta)}\frac{f_k^*(\eta')}{a(\eta')} \tag{101}$$

We list here the explicit formulas obtained substituting in the above equations the field modes functions f_k for ζ in the various scenarios we consider. Due to the presence of positive- and negative-energy solutions, the Whightman function will also have a positive- and negative-energy, which we indicate with $G_k^\pm$.

- *Standard scenario and modified vacuum.* From equations (7), (14)

$$G_k^\pm(0,\eta') = \pm\frac{H^2}{\dot\phi^2}\frac{H^2}{2k^3}\,(1\mp ik\eta')e^{\pm ik\eta'}, \qquad \partial_{\eta'}G_k^\pm(0,\eta') = \pm\frac{H^2}{\dot\phi^2}\frac{H^2}{2k^3}\,k^2\eta' e^{\pm ik\eta'}, \tag{102}$$

- *Modified dispersion relations.* From equation (10), we find that in region IV the Whightman function is the same as the one in equation (102), while in region III it is obtained from (92). Thus,

$$G^\pm_{\mathrm{III},k}(\eta\sim 0,\eta') = \frac{H^2}{\dot\phi^2}\frac{e^{i\frac{\pi}{2}}}{2k^2}\frac{H}{a(\eta')}\chi^{(*)}(k,\eta')\,e^{\pm i\frac{\Lambda}{H}\Omega(k,\eta')}, \quad \chi(k,\eta') = \begin{cases}\frac{1}{\sqrt{U(k,\eta')}} & \text{in region III}\\ (1-\frac{i}{k\eta'}) & \text{in region IV}\end{cases} \tag{103}$$

The derivative of the Whightman function can be written in a form valid both for regions IV and III, given the limiting behaviour of Ω and U at early times, so that

$$\partial_{\eta'}G_k^\pm(\eta\sim 0,\eta') = -\frac{H^2}{\dot\phi^2}\frac{e^{i\frac{\pi}{2}}\,H}{2k}\frac{\gamma^{(*)}(k,\eta')}{a(\eta')}e^{\pm i\frac{\Lambda}{H}\Omega(k,\eta')}, \quad \gamma(k,\eta') = \begin{cases}\frac{1}{\sqrt{U}}\left(\frac{1}{2k}\frac{U'}{U}+iU+\frac{\mathcal{H}}{k}\right) & \text{in region III}\\ i & \text{in region IV}\end{cases} \tag{104}$$

where the suffix $^{(*)}$ means that the positive-energy solution is associated with γ^*,χ^*, while the negative-energy with γ,χ. Finally, $\mathcal{H}$ is the conformal Hubble scale.

A.3 Time integral in bispectrum computations with modified dispersion relations

When computing the bispectrum, we have encountered various Laplace integrals of the form

$$\mathcal{J}_j^{(w)} = \int_{y_{\mathrm{II}}}^{y} dy' y'^{w-1} e^{i\frac{\Lambda}{H}x_1\tilde v_{\theta_j}(y')}\,h(x_1,x_S,\theta_j,y')\,, \tag{105}$$

see equations (35), (53), (70), where w and $h(x_1,x_S,\theta_j,y')$ change in the different integrals and the variables are defined in (34).

In particular, in the case of the minimal coupling interaction (22), the integral present in the correction of order β_k to the bispectrum, see equation (35), has the form of (105) with $w=1$ and $h(\{x\},\theta_j,y')\to g(\{x\},\theta_j,y')$, where [16]

$$g(\{x_{h\neq j}\},x_j,y',m) \equiv \prod_{h\neq j}\gamma^*(x_h,y')\tilde\gamma(x_j,y')\,e^{i\frac{H}{\Lambda}\int^{y'}\left(\sum_{h\neq j}S_2(x_h)+(-1)^m S_2(x_j)\right)} \tag{106}$$

with γ defined in (104) and

$$S_2(x_i,y) = -\frac{\partial_y^2\omega(x_iy)}{4\omega(x_iy)^2} + 3\frac{(\partial_y\omega(x_iy))^2}{8\omega(x_iy)^3} - \frac{1}{\omega(x_iy)y^2}\,, \qquad \tilde\gamma(x,y') = \begin{cases}\gamma^*(x,y') & \text{if } m=0\\ \gamma(x,y') & \text{if } m=1\end{cases}. \tag{107}$$

	$\lvert y\rvert \ll 1$	$y = y_{\rm II}$	$y \simeq y_*$
$F(x_S, y)$	$1+\,(-y)^\kappa F^{(k)}$	$F(-1)$	$F_* + F_*^{(\nu_*)}\,\lvert y-y_*\rvert^{\nu_*}$
$\tilde v_{\theta_j}(y)$	$y\, v_{\theta_j} - (-y)^{\kappa+1} F^{(k)} \cos(\theta_j)$	$-1-\omega(x_S,-1)$	$\tilde v_{\theta_j}(y_*) + \tilde v_{\theta_j}^{(\nu_*)}(y_*)\,\lvert y-y_*\rvert^{\nu_*}$ or $\tilde v_{\theta_j}(y_*) + \tilde v_{\theta_j}^{(\nu_*)}(y_*)$ $\times\lvert y\text{-}y_*\rvert^{\nu_*}\mathrm{sign}(y\text{-}y_*)$
$g(y)$	$-i(1+\lvert y\rvert^\kappa F^{(k)})$	$-iF(-1)$	$(y-y_*)^{\mu_*-1} g^{(\mu_*-1)}(y_*)$
$q(\theta_j, x_1, y)$	$-4i(v_{\theta_j} + 2\lvert y\rvert^\kappa F^{(k)})$ $\text{-}4i(\chi^*(x_1,y)\text{-}1)\cos(\theta_j)$	$-6i\lvert\gamma^*(1,\text{-}1)\rvert^2 + 2iU(1,\text{-}1)^{\text{-}1}$ $\text{-}4i\cos(\theta_j)\chi^*(x_1,\text{-}1)$	$(y-y_*)^{\tau_*} q^{(\tau_*)}(\theta_j, y_*)$

Table 1: Behaviour of functions entering the integrands in equations (35), (53), (70) close to the boundary points (limits of integration) $y_{\rm II} \sim$ -1 and $y \sim$ -$\frac{H}{\Lambda} \ll$ -1, and close to the possible stationary point(s) y_* in the squeezing limit $x_1 \ll 1$, $x_S \simeq 1$.

In the squeezing limit, for $m = 1$ and $j = 2, 3$, looking at (93) and (104), we obtain

$$g(y) \;\equiv\; g(x_1{\ll}1, x_{j=2,3}{\simeq}x_S{\simeq}1, y, 1) \;\simeq\; -i\,\lvert\gamma(x_S, y)\rvert^2\, e^{i\frac{H}{\Lambda}\mathcal{O}(x_1,\frac{H}{\Lambda})} \;\simeq\; -i\lvert\gamma(1,y)\rvert^2 \qquad (108)$$

In the case, instead, of the higher derivative interaction in equation (43), the integral appearing in equation (53) has $w = 3$ and $h(\{x\},\theta_j,y') \to q(\{x\},\theta_j,y')$, where [16]:

$$\begin{aligned} q(\{x_h\},x_j,y) = \Bigg(6\prod_{h\neq j}\gamma^*(x_h)\gamma(x_j) + \frac{2\vec{x}_{j+1}\cdot\vec{x}_{j+2}}{x_{j+1}\,x_{j+2}}\,\chi^*(x_{j+1})\chi^*(x_{j+2})\gamma(x_j) + \frac{2\vec{x}_{j+1}\cdot\vec{x}_j}{x_{j+1}\,x_j}\,\chi^*(x_{j+1})\chi(x_j)\gamma^*(x_{j+2}) \\ + \frac{2\vec{x}_{j+2}\cdot\vec{x}_j}{x_{j+2}\,x_j}\,\chi^*(x_{j+2})\chi(x_j)\gamma^*(x_{j+1})\Bigg)\, e^{i\frac{H}{\Lambda}\int^y\left(\sum_{h\neq j} S_2(x_h) - S_2(x_j)\right)}\,, \qquad (109)\end{aligned}$$

and S_2 has been defined in equation (107) and χ in (103). We write explicitly only the dependence on x of γ, χ to avoid cluttering the formula. The labels of the x's are defined modulo 3, here.

In the squeezing limit, for $j = 2, 3$, looking at (93) and (104),

$$q(\theta_j, x_1, y) \;\equiv q(x_1{\ll}1, x_{j=2,3}{\simeq}x_S{\simeq}1, y) \;\;\simeq\;\; -6i\lvert\gamma^*(1,y)\rvert^2 + 2i\frac{1}{U(1,y)} - 4i\chi^*(x_1,y)\cos(\theta_j) \qquad (110)$$

For $\frac{\Lambda}{H}x_1 \gg 1$, the integrals like (105) are dominated by the contribution of boundary and stationary points, where the oscillations due to the large phase of the integrand are reduced. It is therefore important to study the behavior at leading order of the integrands in (35), (53), (70) close to the boundary points (limits of integration) $y_{\rm II} \sim -1$ and $y \sim$ -$\frac{H}{\Lambda} \ll -1$ and to the possible stationary point(s) $\{y_*\}$. In table 1, we report the leading and next-to-leading behaviour.

In the table, we include also the function $F(x, y')$, defined in (91), and now written in terms of the variables (34). From its definition (91), it appears that the expansion in powers of $\frac{p}{\Lambda} = -y$ for $\lvert y\rvert \ll 1$ is in fact the expansion for the effective frequency from the effective action. In particular, $\kappa{+}1$ is thus the power of the first correction to the standard physical dispersion relation (model dependent):

$$\omega_{\rm phys}(p) \sim p(1 + \left(\tfrac{p}{\Lambda}\right)^\kappa F^{(\kappa)} + \cdots)\,. \qquad (111)$$

In expanding $g(y)$, $q(\theta_j, x_1, y)$ – see (108), (110) – we have taken into account the WKB conditions. The powers ν_*, μ_*, τ_* are model dependent. We wish to be generic, so $\kappa, \nu_*, \mu_*, \tau_*$ can be fractional or integer[21]. Then, since $F = \frac{\omega_{\rm phys}}{p}$ is a real function, its expansion must be clearly in powers of $|y|$, $|y–y_*|$, with real coefficients, for generic ν_*, κ. Similarly, it must be so for $\tilde{v}_{\theta_j}(x, y)\Lambda = -(p + \omega_{\rm phys}(p))$, which is a real function as well (in table 1 we have been very generic concerning its expansion around y_*).

References

[1] E. Komatsu *et al.*, arXiv:0902.4759 [astro-ph.CO]

[2] V. Desjacques and U. Seljak, Class. Quant. Grav. **27** (2010) 124011 [arXiv:1003.5020 [astro-ph.CO]].

[3] N. Dalal, O. Dore, D. Huterer and A. Shirokov, Phys. Rev. D **77**, 123514 (2008) [arXiv:0710.4560 [astro-ph]].

S. Matarrese and L. Verde, Astrophys. J. **677** (2008) L77 [arXiv:0801.4826 [astro-ph]].

A. Slosar, C. Hirata, U. Seljak, S. Ho and N. Padmanabhan, JCAP **0808**, 031 (2008) [arXiv:0805.3580 [astro-ph]].

[4] J. M. Maldacena, JHEP **0305**, 013 (2003) [arXiv:astro-ph/0210603].

[5] P. Creminelli and M. Zaldarriaga, JCAP **0410** (2004) 006 [arXiv:astro-ph/0407059].

[6] P. Creminelli, G. D'Amico, M. Musso and J. Norena, arXiv:1106.1462 [astro-ph.CO].

[7] L. Verde and S. Matarrese, Astrophys. J. **706** (2009) L91 [arXiv:0909.3224 [astro-ph.CO]].

[8] F. Schmidt and M. Kamionkowski, Phys. Rev. D **82** (2010) 103002 [arXiv:1008.0638 [astro-ph.CO]].

[9] P. D. Meerburg, J. P. van der Schaar and P. S. Corasaniti, JCAP **0905**, 018 (2009) [arXiv:0901.4044 [hep-th]].

[10] C. Wagner and L. Verde, arXiv:1102.3229 [astro-ph.CO].

[11] I. Agullo, L. Parker, Phys. Rev. **D83** (2011) 063526. [arXiv:1010.5766 [astro-ph.CO]].

[12] J. Martin and R. H. Brandenberger, Phys. Rev. D **63**, 123501 (2001) [arXiv:hep-th/0005209].

J. Martin and R. H. Brandenberger, Phys. Rev. D **65**, 103514 (2002) [arXiv:hep-th/0201189].

J. Martin and R. Brandenberger, Phys. Rev. D **68** (2003) 063513 [arXiv:hep-th/0305161].

M. Lemoine, M. Lubo, J. Martin and J. P. Uzan, Phys. Rev. D **65** (2002) 023510 [arXiv:hep-th/0109128].

[13] U. H. Danielsson, Phys. Rev. D **66**, 023511 (2002) [arXiv:hep-th/0203198].

U. H. Danielsson, JHEP **0207** (2002) 040 [arXiv:hep-th/0205227].

[14] K. Schalm, G. Shiu and J. P. van der Schaar, JHEP **0404** (2004) 076 [arXiv:hep-th/0401164].

K. Schalm, G. Shiu and J. P. van der Schaar, AIP Conf. Proc. **743** (2005) 362 [arXiv:hep-th/0412288].

---

[21] Given the WKB conditions, it is $\mu_* \geq 1$, while $\tau_* \geq 0$.

[15] R. Holman and A. J. Tolley, JCAP **0805**, 001 (2008) [arXiv:0710.1302 [hep-th]].

[16] D. Chialva, JCAP **1201**, 037 (2012) [arXiv:1106.0040 [hep-th]].

[17] P. Horava, Phys. Rev. **D79**, 084008 (2009). [arXiv:0901.3775 [hep-th]].

[18] D. J. H. Chung and K. Freese, Phys. Rev. D **61** (2000) 023511 [arXiv:hep-ph/9906542].

D. J. H. Chung and K. Freese, Phys. Rev. D **62**, 063513 (2000) [arXiv:hep-ph/9910235].

D. J. H. Chung, E. W. Kolb and A. Riotto, Phys. Rev. D **65**, 083516 (2002) [arXiv:hep-ph/0008126].

C. Csaki, J. Erlich and C. Grojean, Nucl. Phys. B **604**, 312 (2001) [arXiv:hep-th/0012143].

S. L. Dubovsky, JHEP **0201**, 012 (2002) [arXiv:hep-th/0103205].

[19] D. Mattingly, Living Rev. Rel. **8** (2005) 5 [arXiv:gr-qc/0502097].

T. Jacobson, S. Liberati and D. Mattingly, Annals Phys. **321** (2006) 150 [arXiv:astro-ph/0505267].

[20] D. Baumann and D. Green, arXiv:1102.5343 [hep-th].

A. J. Tolley and M. Wyman, Phys. Rev. D **81** (2010) 043502 [arXiv:0910.1853 [hep-th]].

[21] P. Creminelli, JCAP **0310** (2003) 003 [arXiv:astro-ph/0306122].

[22] M. Porrati, Proceedings of 13th International Seminar on High-Energy Physics: Quarks 2004, Pushkinskie Gory, Russia, 24-30 May 2004 [arXiv:hep-th/0409210].

M. Porrati, Phys. Lett. B **596** (2004) 306 [arXiv:hep-th/0402038].

[23] D. Babich, P. Creminelli and M. Zaldarriaga, JCAP **0408**, 009 (2004) [arXiv:astro-ph/0405356].

[24] A. Ashoorioon, D. Chialva, U. Danielsson, JCAP **1106** (2011) 034. [arXiv:1104.2338 [hep-th]].

[25] J. Martin and D. J. Schwarz, Phys. Rev. D **67** (2003) 083512 [arXiv:astro-ph/0210090].

[26] C. M. Bender, S. A. Orszag, “Asymptotic Methods for Scientists and Engeneers”, McGraw Hill, 1978, (Springer 1999).

[27] A. Erdélyi, “Asymptotic Expansions”, Dover (New York) 1956.

[28] S. Weinberg, Phys. Rev. D **77** (2008) 123541 [arXiv:0804.4291 [hep-th]].

[29] C. Cheung, A. L. Fitzpatrick, J. Kaplan and L. Senatore, JCAP **0802** (2008) 021 [arXiv:0709.0295 [hep-th]].